

# VU Research Portal

## Traumatic brain injury in children

Konigs, M.

2016

### **document version**

Publisher's PDF, also known as Version of record

[Link to publication in VU Research Portal](#)

### **citation for published version (APA)**

Konigs, M. (2016). *Traumatic brain injury in children: Impact on Brain Structure, Neurocognition & Behavior*. [, Vrije Universiteit Amsterdam].

### **General rights**

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

### **Take down policy**

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

### **E-mail address:**

[vuresearchportal.ub@vu.nl](mailto:vuresearchportal.ub@vu.nl)



# CHAPTER 3

## INTELLIGENCE AFTER TRAUMATIC BRAIN INJURY: META-ANALYSIS OF OUTCOMES AND PROGNOSIS

*Published as:*

Königs, M., Engenhorst, P. J., & Oosterlaan, J. (2016). Intelligence after traumatic brain injury: meta-analysis of outcomes and prognosis. *European journal of neurology*, 23(1), 21-29.

## ABSTRACT

### BACKGROUND

Worldwide, 54-60 million individuals sustain traumatic brain injury (TBI) each year. This meta-analysis aimed to quantify intelligence impairments after TBI, and to determine the value of age and injury severity in the prognosis of TBI.

### METHODS

Electronic database search identified 81 relevant peer-reviewed articles encompassing 3,890 patients. Full scale IQ (FSIQ), performance IQ (PIQ) and verbal IQ (VIQ) impairments were quantified (Cohen's *d*) for patients with mild, moderate and severe TBI in the subacute phase of recovery and the chronic phase. Meta-regressions explored prognostic values of age and injury severity measures for intelligence impairments.

### RESULTS

In the subacute phase, FSIQ impairments were absent for patients with mild TBI, medium-sized for patients with moderate TBI ( $p < .001$ ,  $d = -0.61$ ) and large for patients with severe TBI ( $p < .001$ ,  $d = -1.09$ ). In the chronic phase, FSIQ impairments were small for patients with mild or moderate TBI ( $ps \leq .008$ ,  $ds = -0.37$  and  $-0.19$ ) and large for patients with severe TBI ( $p < .001$ ,  $d = -0.80$ ). Adults with mild TBI had larger PIQ and VIQ impairments in the chronic phase than children ( $Qs \geq 5.21$ ,  $ps \leq .02$ ), while children with severe TBI had larger FSIQ and VIQ impairments than adults ( $Qs \geq 4.40$ ,  $ps \leq .04$ ). GCS score, LOC duration and PTA duration moderately to strongly predicted FSIQ, PIQ and VIQ impairments ( $.41 \leq rs \leq .82$ ,  $ps \leq .02$ ), but no differences in predictive value were observed.

### CONCLUSIONS

TBI causes persisting intelligence impairments, where children may have better recovery from mild TBI and poorer recovery from severe TBI than adults. Injury severity measures predict intelligence impairments and do not outperform one another.

## INTRODUCTION

Yearly, 54-60 million individuals sustain traumatic brain injury (TBI) worldwide (Feigin et al., 2013). Survivors of TBI often have impaired neurocognitive functions, prominently manifested in intelligence. In children, the impact of TBI is thought to slow down post-injury development of intelligence, which may translate into increasing intellectual delays (Crowe, Catroppa, Babl, Rosenfeld, & Anderson, 2012). Intelligence impairments in children with TBI have in turn been associated with psychiatric symptoms and poor academic attainment (Donders & Warschawsky, 2007; Donders, 1994; Thaler et al., 2010). In adults with TBI, intelligence impairments relate to psychopathology, poor vocational placement and lower quality of life (Bowman, 1996; Yasuda, Wehman, Targett, Cifu, & West, 2001). These findings highlight intelligence as a crucial factor in the prognosis of TBI.

Clinicians use the Glasgow Coma Scale (GCS), duration of loss of consciousness (LOC) and/or post-traumatic amnesia (PTA) duration as predictive tools in the prognosis of TBI (Mandleberg, 1976; Winogron, Knights, & Bawden, 1984), while their concurrent prognostic utility remains unclear. Although the clinical application of PTA duration has been held back by marked variability in its definition and measurement methodology (Ahmed, Bierley, Sheikh, & Date, 2000), literature suggests that PTA duration may be superior to the GCS score and LOC duration in predicting a range of outcomes following TBI, including intelligence and functional independence (Greenwood, 1997; Perrin et al., 2014; Sherer, Struchen, Yablon, Wang, & Nick, 2008). These findings warrant the comparison of major prognostic factors for intelligence outcome of TBI. This study is the first to comprehensively quantify the impact of mild, moderate and severe TBI on intelligence in children and adults, and to determine the prognostic value of age at injury and injury severity using meta-analytic methods.


## METHODS

### STUDY SELECTION

This study was performed according to the PRISMA guidelines (Liberati et al., 2009). Included were studies that: (1) investigated patients with TBI; (2) reported on post-injury intelligence using any version of the most-widely used measure of intelligence: the Wechsler Intelligence Scales; (3) reported statistics allowing the calculation of effect sizes; and (4) reported on injury severity according to the mean or median GSC score, LOC duration and/or PTA duration. Excluded were studies that: (1) investigated patients with substance intoxication or prior head injury; (2) reported on injury severity based on < 75% of the patient sample; or (3) reported case studies.

The databases PubMed, Embase, PsycINFO and Web of Knowledge were searched for articles potentially meeting inclusion criteria using combinations of the following standard and thesaurus search terms: 'Wechsler', 'Head', 'Brain', 'Injury' and 'Trauma'. A total of 2,373 records were identified corresponding to 1,125 unique articles published before April 2014 (Figure 1). Titles and abstracts were assessed to further select articles meeting inclusion criteria. Two authors (M.K. and P.E.) independently assessed the full-text articles for eligibility. A total of 117 articles met all inclusion criteria, of which 36 were excluded for possibly overlapping samples based on the period and area of recruitment. Reference lists of included articles were searched for additional articles satisfying the inclusion criteria. Finally, eighty-one articles were included in the current analysis representing 3,890 patients with TBI (see Supporting Information).





**Figure 1.** Flow diagram of the meta-analytic search and study selection  
*Note.* IQ = intelligence quotient.

## MODERATOR AND DEPENDENT VARIABLES

Samples of patients with TBI and controls were identified in the included articles. For these samples, information on injury severity, time post-injury (TPI), demographics and study quality was extracted.

### INTELLIGENCE

Intelligence was defined and measured according age-corrected standard scores ( $M = 100$ ,  $SD = 15$ ) on the Wechsler Preschool and Primary Scale of Intelligence (Wechsler, 1967, 1989), Wechsler Intelligence Scales for Children (Wechsler, 1955, 1974, 1991, 2003) or the Wechsler Adult Intelligence Scales (Wechsler, 1955, 1981, 1997, 2008). Both full and short forms were included, as short forms are known to correlate strongly ( $.76 < r_s \leq .91$ ) with full administrations of the Wechsler scales (Guilmette, Dabrowski, Kennedy, & Gnys, 1999; Sattler, 2001). The included articles allowed extraction of mean full-scale IQ (FSIQ), performance IQ (PIQ) and/or verbal IQ (VIQ) and the corresponding SD for samples of patients with TBI and control participants, if available. Only two studies used the WISC-IV. Therefore meta-analysis of WISC-IV factor scores dissecting FSIQ into verbal comprehension, perceptual reasoning, working memory and processing speed capabilities was not feasible.

### INJURY SEVERITY

One or more of the following measures defined mild, moderate or severe TBI: GCS score (15-13; 12-9; and 8-3), LOC duration (< 30 minutes; 30 minutes - 24 hours; and > 24 hours) and PTA duration (<1 day; 1 - 7 days; and > 7 days, respectively; Tsao, 2010). The definition and measurement methodology of injury severity varied considerably among studies and are described in detail in the Supporting Information.

### TIME POST-INJURY

To allow comparison of short- and long-term predictive values of injury severity measures, we distinguished between reports in the subacute phase of recovery ( $TPI < 6$  months) and the chronic phase ( $TPI \geq 6$  months), in line with existing literature (Babikian & Asarnow, 2009; Königs, de Kieviet, & Oosterlaan, 2012).

### DEMOGRAPHIC VARIABLES

Group means of the demographic variables age, age at injury, years of education and gender were extracted from the articles. Based on age at injury, we discriminated between childhood TBI ( $\leq 18$  years) and adult TBI ( $> 18$  years).



## STUDY QUALITY

The quality of the included studies was independently assessed by two authors (M.K. and P.E.) using the Newcastle Ottawa Scale (NOS, (Wells et al., 2012). The NOS measures study quality concerning subject selection, group comparability and exposure to testing procedures. Higher scores reflect higher study quality (0 - 9 points). Discrepancies between author ratings were resolved by consensus, while inter-rater agreement was adequate (intraclass correlation = .69).

## STATISTICAL ANALYSIS

Statistical analyses were performed using SPSS version 20.0 (SPSS inc., 2013) and Comprehensive Meta-Analysis software (Borenstein, 2005). First, we assessed comparability of the mild, moderate and severe TBI groups within the TPI groups (subacute and chronic phase) and between TPI groups using independent t-tests comparing age, age at injury, gender, years of education, TPI, GCS score (because this measure of injury severity was available for most of the TBI samples) and NOS study quality score.

Second, we determined the intelligence impairment for each sample of TBI patients derived from the included articles. We calculated the effect size (Cohen's *d*) using the mean IQ score and the corresponding SD of the TBI group and the mean IQ score and the corresponding SD of either: (1) the control group for controlled studies; or (2) the Wechsler normative mean and SD for uncontrolled studies. We investigated the validity of using the normative mean for uncontrolled studies by calculating the individual effect sizes for controlled studies using both methods and subsequently comparing the two acquired meta-analytic effect sizes using Q-testing. Effect sizes calculated using the normative mean were somewhat smaller than those calculated using the original control groups for FSIQ ( $\Delta d = 0.17$ ), PIQ ( $\Delta d = 0.10$ ) and VIQ ( $\Delta d = .15$ ), although the differences were not significant ( $ps \geq .12$ ). This indicates that using the normative mean for effect size calculation may provide conservative estimates of IQ impairments following TBI.

Meta-analytic effect sizes were calculated for IQ indices in the TBI severity groups, discriminating between the subacute phase of recovery and the chronic phase. The derived effect sizes were weighted by their inverse variance, thereby accounting for sample size and error of measurement (Borenstein, Hedges, Higgins, & Rothstein, 2009). All meta-analytic effect sizes were calculated using the random model, since heterogeneity may have been introduced by using data from different versions of the Wechsler scales. Heterogeneity of effect sizes was assessed using the  $I^2$  statistic, where values of 25, 50 and 75% were indicative of low, moderate and high heterogeneity, respectively (Higgins, Thompson, Deeks, & Altman, 2003). To track the recovery of intelligence following TBI,

the meta-analytic effect sizes were compared between the TPI groups using *Q*-testing. All meta-analytic effect sizes were interpreted according to Cohen, translating  $d = -0.20$  into small,  $d = -0.50$  as medium and  $d = -0.80$  as large effects (Cohen, 1988).

Subsequently, we investigated the predictive value of age at injury and the selected injury severity measures by performing meta-regressions between the study samples' individual effect sizes for FSIQ, PIQ and VIQ and the study samples' mean age at injury, mean GCS score, mean LOC duration and/or mean PTA duration for the two TPI groups separately. The average values of injury severity measures were rarely reported for patients with mild TBI. To allow inclusion of samples of patients with mild TBI in the meta-regression, we replaced missing values with the median of the mild TBI range for the GCS score, LOC duration and PTA duration (14, 0 days and 0.5 days, respectively; Carroll, Cassidy, Holm, Kraus, & Coronado, 2004). Meta-regression analysis was performed using the random model (method of moments; Borenstein et al., 2009). The predictive values of the GCS score, LOC duration and PTA duration for FSIQ, PIQ and VIQ impairments in the subacute phase of recovery and the chronic phase, were compared using *t*-tests while adjusting *p*-values for multiple comparisons according to Bonferroni (reported  $p = \text{obtained } p * m$ , where *m* denotes the number of comparisons). In all analyses, standardized correlations were interpreted according to Cohen, translating  $r = .10$ ,  $r = .30$  and  $r = .50$  into weak, moderate and strong relations (Cohen, 1988).

The robustness of effect sizes was assessed by calculating Rosenthal's fail-safe *n* to determine the number of studies with a null effect that would be necessary to cancel out significant effect sizes (Rosenthal, 1995). Fail-safe *n* values  $> 5k + 10$  were considered robust, where *k* refers to the number of samples on which the relevant effect size was calculated (Rosenthal, 1995). Publication bias was assessed by Egger Funnel plot asymmetry (Egger, Smith, Schneider, & Minder, 1997). The possibility of a disproportionate influence of small-sized samples or poor quality studies with large effect sizes was investigated, which would become evident by positive associations between effect size and sample size or study quality, respectively. All tests of significance were two-sided with  $\alpha = .05$ .

## RESULTS

### GROUP COMPARABILITY

A description of analyses investigating comparability of the TBI severity groups and TPI groups in terms of age, age at injury, gender ratio, years of education, TPI, GCS score and NOS study quality score is provided in the Supporting Information (see Table 1 in Supporting Information for patient sample characteristics). No significant differences between groups were observed that would meaningfully confound the interpretation of effect sizes. Nevertheless, a large but non-significant difference in PTA duration was identified between the severe TBI groups in the subacute phase ( $M = 34.7$ ,  $SD = 28.9$ ) and chronic phase ( $M = 52.8$ ,  $SD = 33.4$ ). We matched these groups on injury severity during analyses comparing TPI groups, by excluding four studies from the chronic phase group (references 24, 64, 69-70 in Supporting Information), in order to prevent effects of injury severity to confound the analysis of recovery.

### THE EFFECT OF TBI ON INTELLIGENCE

#### MILD TBI

Effect sizes for FSIQ, PIQ and VIQ in the subacute phase of recovery and the chronic phase showed moderate to high heterogeneity (Table 2), indicating that there was considerable variability in the magnitude of intelligence impairments. Patients with mild TBI had no significant intelligence impairments in the subacute phase of recovery, but there were found significant small-sized FSIQ and VIQ impairments in the chronic phase. There were no significant differences between the subacute phase of recovery and the chronic phase in the effects of mild TBI on any of the IQ indices ( $Qs(1) \leq 0.5$ ,  $ps \geq .49$ ), and the VIQ impairment in the chronic phase did not differ from the PIQ impairment ( $Q(1) = 0.7$ ,  $p = .41$ ).

#### MODERATE TBI

Effect sizes reflecting the effect of moderate TBI on intelligence showed low heterogeneity. Patients with moderate TBI had significant medium-sized FSIQ and PIQ impairments and a small VIQ impairment in the subacute phase of recovery. In the chronic phase, there were found significant small-sized FSIQ and VIQ impairments, but no impairment in PIQ was observed. The FSIQ and PIQ impairments were smaller in the chronic phase as compared to the subacute phase ( $Q(1) = 7.6$ ,  $p = .006$  and  $Q(1) = 10.4$ ,  $p = .001$ , respectively), indicating ameliorating effects of recovery. There was no difference in the magnitude of VIQ impairments between the subacute and chronic phase ( $Q(1) = 0.9$ ,  $p = .33$ ). No significant difference was observed between the magnitude of PIQ and VIQ impairments in the subacute phase of recovery ( $Q(1) = 1.9$ ,  $p = .17$ ) or the chronic phase ( $Q(1) = 1.4$ ,  $p = 0.23$ ).

## SEVERE TBI

There was found moderate to high heterogeneity in meta-analytic effect sizes reflecting the impact of severe TBI on intelligence. Patients with severe TBI exhibited significant large-sized FSIQ, PIQ and VIQ impairments in the subacute phase of recovery. In the chronic phase, severe TBI patients showed large-sized FSIQ and PIQ impairments, and a medium-sized VIQ impairment. There were found recovery effects for FSIQ and PIQ impairments ( $Qs \geq 3.9$ ,  $ps \leq .05$ ), but not for VIQ impairments ( $Q(1) = 1.9$ ,  $p = .17$ ). Severe TBI patients had significantly larger PIQ impairments as compared to VIQ impairments in the subacute phase of recovery ( $Q(1) = 6.7$ ,  $p = .01$ ), but such a difference did not reach conventional threshold of significance in the chronic phase ( $Q(1) = 3.1$ ,  $p = .08$ ).

**Table 2.** Meta-analytic effect sizes for intelligence impairments of patients with mild, moderate and severe TBI


| Group          | FSIQ |       |                |       |                |      |       |      |        | PIQ |       |                |       |                |      |       |      |        |
|----------------|------|-------|----------------|-------|----------------|------|-------|------|--------|-----|-------|----------------|-------|----------------|------|-------|------|--------|
|                | k    | ES    | 95% CI         | p     | I <sup>2</sup> | fsn  | p(EF) | p(n) | p(NOS) | k   | ES    | 95% CI         | p     | I <sup>2</sup> | fsn  | p(EF) | p(n) | p(NOS) |
| Mild TBI       |      |       |                |       |                |      |       |      |        |     |       |                |       |                |      |       |      |        |
| Subacute Phase | 10   | -0.29 | -0.64 to 0.06  | .11   | 78%            | 0    | .56   | .95  | .28    | 10  | -0.10 | -0.36 to 0.17  | .47   | 70%            | 0    | .69   | .79  | .41    |
| Chronic Phase  | 16   | -0.37 | -0.58 to -0.15 | <.001 | 63%            | 119  | .65   | .67  | .12    | 15  | -0.16 | -0.44 to 0.13  | .29   | 83%            | 0    | .79   | .89  | .18    |
| Moderate TBI   |      |       |                |       |                |      |       |      |        |     |       |                |       |                |      |       |      |        |
| Subacute Phase | 5    | -0.61 | -0.89 to -0.34 | <.001 | 0%             | 20   | .55   | .17  | .11    | 6   | -0.72 | -1.02 to -0.42 | <.001 | 15%            | 28   | .30   | .44  | .68    |
| Chronic Phase  | 11   | -0.19 | -0.32 to -0.05 | .008  | 0%             | 0    | .09   | .18  | .75    | 9   | -0.10 | -0.31 to 0.12  | .12   | 32%            | 0    | .12   | .23  | .74    |
| Severe TBI     |      |       |                |       |                |      |       |      |        |     |       |                |       |                |      |       |      |        |
| Subacute Phase | 21   | -1.09 | -1.34 to -0.83 | <.001 | 75%            | 4767 | .78   | .66  | .09    | 21  | -1.21 | -1.45 to -0.97 | <.001 | 69%            | 1602 | .27   | .84  | .48    |
| Chronic Phase  | 40   | -0.80 | -0.97 to -0.63 | <.001 | 76%            | 3647 | .50   | .33  | .44    | 29  | -0.89 | -1.07 to -0.70 | <.001 | 72%            | 2188 | .46   | .97  | .14    |
| VIQ            |      |       |                |       |                |      |       |      |        |     |       |                |       |                |      |       |      |        |
| Group          | k    | ES    | 95% CI         | p     | I <sup>2</sup> | fsn  | p(EF) | p(n) | p(NOS) |     |       |                |       |                |      |       |      |        |
| Mild TBI       |      |       |                |       |                |      |       |      |        |     |       |                |       |                |      |       |      |        |
| Subacute Phase | 9    | -0.19 | -0.43 to 0.05  | .13   | 55%            | 0    | .18   | .23  | .41    |     |       |                |       |                |      |       |      |        |
| Chronic Phase  | 15   | -0.30 | -0.50 to -0.10 | .003  | 58%            | 67   | .47   | .69  | .26    |     |       |                |       |                |      |       |      |        |
| Moderate TBI   |      |       |                |       |                |      |       |      |        |     |       |                |       |                |      |       |      |        |
| Subacute Phase | 6    | -0.45 | -0.71 to -0.19 | <.001 | 0%             | 7    | .40   | .98  | .22    |     |       |                |       |                |      |       |      |        |
| Chronic Phase  | 9    | -0.30 | -0.46 to -0.13 | <.001 | 0%             | 9    | .05   | .08  | .40    |     |       |                |       |                |      |       |      |        |
| Severe TBI     |      |       |                |       |                |      |       |      |        |     |       |                |       |                |      |       |      |        |
| Subacute Phase | 21   | -0.80 | -0.98 to -0.63 | <.001 | 44%            | 813  | .56   | .88  | .53    |     |       |                |       |                |      |       |      |        |
| Chronic Phase  | 31   | -0.66 | -0.83 to -0.48 | <.001 | 71%            | 1475 | .72   | .46  | .06    |     |       |                |       |                |      |       |      |        |

Note. Significant effect sizes are displayed in bold. CI = confidence interval; ES = effect size (Cohen's *d*); FSIQ = full-scale intelligence quotient; fsn = fail-safe *n*; *k* = number of samples; PIQ = performance intelligence quotient; *p*(EF) = *p* value of Egger funnel plot asymmetry; *p*(*n*) = *p* value of the relation between sample size and effect size; *p*(NOS) = *p* value of the relation between Newcastle Ottawa Scale quality score and effect size; TBI = traumatic brain injury; VIQ = verbal intelligence quotient.

### THE ROLE OF AGE AT INJURY

Meta-regressions between age at injury and FSIQ, PIQ or VIQ impairments did not reach significance in the subacute or in the chronic phase of recovery ( $ps \geq .14$ ). To rule out effects of age at injury, we also explored group differences between child and adult TBI groups (Figure 2).

Children and adults with mild TBI showed no differences in FSIQ, PIQ and VIQ impairments in the subacute phase ( $Qs \leq .33, ps \geq .56$ ). In the chronic phase, no difference was observed in FSIQ ( $Q(1) = 3.7, p = .06$ ), but adults with mild TBI had larger PIQ and VIQ impairments than children ( $Qs \geq 5.2, ps \leq .02$ ). No differences were found between children and adults with moderate TBI ( $Qs \leq 2.8, ps \geq .09$ ) and between children and adults with severe TBI in the subacute phase ( $Qs \leq 1.3, ps \geq .25$ ). In contrast, children with severe TBI in the chronic phase had larger FSIQ and VIQ impairments than adults ( $Qs \geq 4.4, ps \leq .04$ ), while no difference was observed in PIQ ( $Q(1) = 0.5, p = .50$ ).



**Figure 2.** Group comparisons between children and adults with mild, moderate and severe TBI

Note. FSIQ = full-scale intelligence quotient; k = number of samples; TBI = traumatic brain injury; PIQ = performance intelligence quotient; VIQ = verbal intelligence quotient.

The observed differences between children and adults with mild TBI could not be explained by differential recovery tracks, as we found no recovery of IQ after mild TBI in children as well as adults ( $Qs \leq 3.3$ ,  $ps \geq .07$ ). In contrast, adults with severe TBI showed significant recovery of FSIQ, PIQ and VIQ impairments ( $Qs \geq 6.7$ ,  $ps \leq .01$ ), while no recovery was observed in children with severe TBI ( $Qs \leq 0.4$ ,  $ps \geq .54$ ). The observed differences between children and adults, and between the subacute and chronic phase of recovery, were not confounded by differences in GCS score, LOC duration or PTA duration ( $ts \leq 1.4$ ,  $ps \geq .21$ ).

### PREDICTING INTELLIGENCE IMPAIRMENTS USING GCS, LOC AND PTA

#### GCS SCORE

Meta-regression analysis revealed that lower GCS scores strongly predicted larger FSIQ, PIQ and VIQ impairments among patients with mild to severe TBI in the subacute phase of recovery (see Table 3). In the chronic phase, GCS scores were not predictive of FSIQ impairments, but lower GCS scores were moderately predictive of larger PIQ and VIQ impairments.

#### LOC DURATION

Visual inspection of the meta-regression plots revealed two multivariate outliers that were removed from the analyses (references 30 and 66 in Supporting Information). Longer LOC duration strongly predicted larger FSIQ, PIQ and VIQ impairments in the subacute phase of recovery. In the chronic phase, longer LOC duration strongly predicted larger FSIQ and PIQ impairments and moderately predicted larger VIQ impairments.

**Table 3.** Predictive values of GCS scores, LOC duration and PTA duration for intelligence

|                       | GCS score |          |      |                 | LOC duration |          |      |                 | PTA duration |          |      |                 |
|-----------------------|-----------|----------|------|-----------------|--------------|----------|------|-----------------|--------------|----------|------|-----------------|
|                       | <i>k</i>  | <i>r</i> | SE   | <i>p</i>        | <i>k</i>     | <i>r</i> | SE   | <i>p</i>        | <i>k</i>     | <i>r</i> | SE   | <i>p</i>        |
| <i>Subacute phase</i> |           |          |      |                 |              |          |      |                 |              |          |      |                 |
| FSIQ                  | 23        | .63      | 0.14 | <b>.001</b>     | 22           | -.73     | 0.11 | <b>&lt;.001</b> | 20           | -.69     | 0.13 | <b>&lt;.001</b> |
| PIQ                   | 28        | .69      | 0.11 | <b>&lt;.001</b> | 18           | -.82     | 0.08 | <b>&lt;.001</b> | 24           | -.76     | 0.09 | <b>&lt;.001</b> |
| VIQ                   | 27        | .54      | 0.14 | <b>.004</b>     | 18           | -.76     | 0.11 | <b>&lt;.001</b> | 24           | -.58     | 0.15 | <b>.003</b>     |
| <i>Chronic phase</i>  |           |          |      |                 |              |          |      |                 |              |          |      |                 |
| FSIQ                  | 43        | .22      | 0.15 | .15             | 39           | -.64     | 0.10 | <b>&lt;.001</b> | 28           | -.36     | 0.17 | .06             |
| PIQ                   | 29        | .43      | 0.16 | <b>.02</b>      | 35           | -.58     | 0.12 | <b>&lt;.001</b> | 21           | -.64     | 0.14 | <b>.002</b>     |
| VIQ                   | 31        | .41      | 0.16 | <b>.004</b>     | 36           | -.44     | 0.14 | <b>.008</b>     | 22           | -.52     | 0.17 | <b>.01</b>      |

*Note.* Significant effect sizes are displayed in bold. FSIQ = full-scale intelligence quotient; GCS = Glasgow Coma Scale; *k* = number of samples; LOC = loss of consciousness; PIQ = performance intelligence quotient; PTA = post-traumatic amnesia; VIQ = verbal intelligence quotient; SE = standard error.



### PTA DURATION

Longer PTA duration strongly predicted larger FSIQ, PIQ and VIQ impairments in the subacute phase of recovery. In the chronic phase, longer PTA duration was moderately predictive of larger FSIQ impairments, although this effect just escaped conventional levels of significance ( $p = .06$ ), whereas longer PTA duration strongly predicted larger PIQ and VIQ impairments. Importantly, there were no significant differences between the predictive values of the GCS score, LOC duration and PTA duration for FSIQ, PIQ and VIQ impairments in the subacute phase of recovery ( $ts \leq 1.1$ , adjusted  $ps \geq .99$ ) and the chronic phase ( $ts \leq 2.3$ , adjusted  $ps \geq .36$ ).

### PUBLICATION BIAS

An elaborate description of publication bias analysis (Table 2) is provided in the Supporting Information, indicating that it is unlikely that publication bias meaningfully influenced the reported effect sizes for the mild, moderate and severe TBI groups, with the single exception of VIQ impairments in the chronic phase after moderate TBI.

## DISCUSSION

To our best knowledge, current meta-analysis of 81 peer-reviewed articles encompassing 3,890 patients with TBI is the most comprehensive study quantifying the impact of TBI on intelligence, and the first to use meta-analytic methods to elucidate the role of age at injury and compare the predictive powers of major prognostic factors for the magnitude of intelligence deficits after TBI. The results show that both moderate and severe TBI cause persisting intelligence impairments. Children seem to have better long-term outcome of mild TBI and poorer long-term outcome of severe TBI than adults. The GCS score, LOC duration and PTA duration moderately to strongly predict intelligence impairments of patients with TBI, but these prognostic factors do not outperform one another in predicting these impairments.

Patients with mild TBI showed no meaningful intelligence impairments in the subacute phase of recovery, but there were found small impairments in FSIQ and VIQ in the chronic phase. Patients with moderate TBI exhibited medium-sized FSIQ and PIQ impairments and small VIQ impairments in the subacute phase of recovery. The medium-sized FSIQ impairment as observed in the subacute phase reduced to a small effect in the chronic phase, whereas the medium-sized effect on PIQ as observed in the subacute phase disappeared in the chronic phase. No change was observed for the VIQ impairment in patients with moderate TBI. Patients with severe TBI exhibited large-sized FSIQ, PIQ and VIQ impairments in the subacute phase of recovery that persisted into the chronic phase, although the VIQ impairment did no longer pass the threshold for a large effect and was medium-sized in the chronic phase.

Patients with severe TBI showed stronger depressions of PIQ as compared to VIQ in the subacute phase, indicating that fluid aspects of intelligence (speed of information processing, attention and visuomotor functioning) are more vulnerable to the impact of TBI than crystallized aspects of intelligence (verbal knowledge). Recent studies using the latest Wechsler Intelligence Scale indicate that PIQ impairments after TBI are mainly accounted for by impairments in speed of processing (Donders & Strong, 2014). The discrepancy between PIQ and VIQ impairments in patients with severe TBI was not found in the chronic phase, likely reflecting relative recovery of processing speed over time.

Analyses differentiating between childhood and adult TBI suggest that the long-term outcome of mild TBI may be more favorable for children than adults. In contrast, children with severe TBI in the chronic phase had larger FSIQ and VIQ impairments than adults, traced back to an absence of recovery in children relative to distinct recovery of FSIQ, PIQ and VIQ impairments in adults. In line with the 'growing into deficit' theory, PIQ impairments in children with severe TBI may slow post-injury development of VIQ, in turn causing progressive delays in verbal skills.

The included studies applied highly variable definitions of injury severity and measurement

methodology. It was found that there is no consensus on the most suitable type of GCS score to be reported, the definition of LOC resolution, the definition of PTA duration and PTA measurement. Given the wide range of definitions and measures used to assess injury severity, it is striking that many studies do not report the definitions and measures applied. The results of the current meta-analysis show that lower GCS scores, longer LOC duration and longer PTA duration strongly predicted greater FSIQ, PIQ and VIQ impairments in the subacute phase. In the chronic phase, longer LOC duration moderately to strongly predicted FSIQ, PIQ and VIQ impairments, while lower GCS scores and longer PTA duration moderately to strongly predicted larger PIQ and VIQ impairments. Longer PTA duration showed moderate predictive power for larger FSIQ impairments in the chronic phase, but this effect just escaped the conventional level of significance. There was no evidence for differences between the GCS score, LOC duration and PTA duration regarding their predictive power for intelligence outcome of patients with mild to severe TBI, contradicting literature suggesting that PTA duration is superior in predicting intelligence after TBI.

This study has some limitations. First, we included uncontrolled studies and used the normative mean to calculate effect sizes for these studies. However, we also showed that this strategy to maximize study inclusion did not meaningfully influence effect sizes and would otherwise have provided slightly conservative estimates. Second, we applied a cross-sectional approach to track recovery effects of time, instead of including only (the scarcely available) longitudinal studies. However, for all comparisons we showed that the subacute and chronic groups involved were comparable in terms of demographic information and injury parameters. Third, some comparisons between childhood and adult TBI involved limited numbers of studies and should therefore be interpreted with caution. Fourth, included studies provided insufficient information to investigate the effect of injury mechanism (e.g. open vs. closed TBI). Last, it is likely that heterogeneity in the definition and measurement of injury severity contributed to measurement error in meta-regression analyses.

In conclusion, the current meta-analysis on a very large sample of patients with TBI indicates that intelligence impairments persist after moderate to severe TBI, while children may have better long-term outcome of mild TBI and poorer long-term outcome from severe TBI than adults. Although injury severity measures were shown to powerfully predict the future of patients with TBI, this study did not yield evidence for superiority of one or more of the measures in predicting intelligence outcome. We advise clinicians to recognize the importance of age at injury and the use of standardized measures of injury severity in the prognosis of intelligence after TBI.

## REFERENCES

- Ahmed, S., Bierley, R., Sheikh, J. I., & Date, E. S. (2000). Review of subject post-traumatic amnesia after closed head injury: a review of the literature and some suggestions for further research. *Brain Injury*, 14(9), 765–780.
- Babikian, T., & Asarnow, R. (2009). Neurocognitive outcomes and recovery after pediatric TBI: meta-analytic review of the literature. *Neuropsychology*, 23(3), 283–296. doi:10.1037/a0015268
- Borenstein, M. (2005). *Comprehensive meta-analysis*. Englewood, NJ: Biostat.
- Borenstein, M., Hedges, L. V., Higgins, J. P. T., & Rothstein, H. R. (2009). *Introduction to Meta-Analysis*. West Sussex: John Wiley & Sons, Ltd.
- Bowman, M. L. M. (1996). Ecological validity of neuropsychological and other predictors following head injury. *The Clinical Neuropsychologist*, 10(4), 382–396. doi:10.1080/13854049608406699
- Carroll, L., Cassidy, J. D., Holm, L., Kraus, J., & Coronado, V. (2004). Methodological issues and research recommendations for mild traumatic brain injury: the who collaborating center task force on mild traumatic brain injury. *Journal of Rehabilitation Medicine*, 36(2), 113–125. doi:10.1080/16501960410023877
- Cohen, J. (1988). *Statistical power analysis for the behavioral science*. Hillsdale, NY: Erlbaum.
- Crowe, L. M., Catroppa, C., Babl, F. E., Rosenfeld, J. V., & Anderson, V. A. (2012). Timing of traumatic brain injury in childhood and intellectual outcome. *Journal of Pediatric Psychology*, 37(7), 745–754. doi:10.1093/jpepsy/jps070
- Donders, J. (1994). Academic placement after traumatic brain injury. *Journal of School Psychology*, 32(1), 53–65. doi:10.1016/0022-4405(94)90028-0
- Donders, J., & Strong, C.-A. H. (2014). Clinical Utility of the Wechsler Adult Intelligence Scale-Fourth Edition After Traumatic Brain Injury. *Assessment*, 22(1), 17–22.
- Donders, J., & Warschawsky, S. (2007). Neurobehavioral outcomes after early versus late childhood traumatic brain injury. *The Journal of Head Trauma Rehabilitation*, 22(5), 296–302. doi:10.1097/01.HTR.0000290974.01872.82
- Egger, M., Smith, G. D., Schneider, M., & Minder, C. (1997). Bias in meta-analysis detected by a simple graphical test. *BMJ*, 315, 629–634.
- Feigin, V. L., Theadom, A., Barker-Collo, S., Starkey, N. J., McPherson, K., Kahan, M., ... Ameratunga, S. (2013). Incidence of traumatic brain injury in New Zealand: a population-based study. *Lancet Neurology*, 12(1), 53–64. doi:10.1016/S1474-4422(12)70262-4
- Greenwood, R. (1997). Value of recording duration of post-traumatic amnesia. *Lancet*, 349, 1041–1042.
- Guilmette, T. J., Dabrowski, J., Kennedy, M. L., & Gnys, J. (1999). A Comparison of Nine WAIS-R Short Forms in Individuals with Mild to Severe Traumatic Brain Injury. *Assessment*, 6(1), 33–41. doi:10.1177/107319119900600104
- Higgins, J. P. T., Thompson, S. G., Deeks, J. J., & Altman, D. G. (2003). Measuring inconsistency in meta-analyses. *BMJ (Clinical Research Ed.)*, 327(7414), 557–60. doi:10.1136/bmj.327.7414.557
- Königs, M., de Kieviet, J. F., & Oosterlaan, J. (2012). Post-traumatic amnesia predicts intelligence impairment following traumatic brain injury: a meta-analysis. *Journal of Neurology, Neurosurgery, and Psychiatry*, 83(11), 1048–1055. doi:10.1136/jnnp-2012-302635
- Liberati, A., Altman, D. G., Tetzlaff, J., Mulrow, C., Gøtzsche, P. C., Ioannidis, J. P. a, ... Moher, D. (2009). The PRISMA statement for reporting systematic reviews and meta-analyses of studies that evaluate health care interventions: explanation and elaboration. *PLoS Medicine*, 6(7), 1–28.
- Mandleberg, I. A. (1976). Cognitive recovery after severe head injury. 3. WAIS verbal and performance IQs as a function of post-traumatic amnesia duration and time from injury. *Journal of Neurology, Neurosurgery & Psychiatry*, 39(10), 1001–1007.

- Perrin, P. B., Niemeier, J. P., Mougeot, J.-L., Vannoy, C. H., Hirsch, M. A., Watts, J. A., ... Whitney, M. P. (2014). Measures of Injury Severity and Prediction of Acute Traumatic Brain Injury Outcomes. *The Journal of Head Trauma Rehabilitation*, 30(2):136-42.
- Rosenthal, R. (1995). Writing meta-analytic reviews. *Psychological Bulletin*, 118(2):183-192. doi:10.1037/0033-2909.118.2.183
- Sattler, J. M. (2001). Assessment of children: cognitive applications. Jerome M. Sattler, *San Diego*, 774.
- Sherer, M., Struchen, M. A., Yablon, S. A., Wang, Y., & Nick, T. G. (2008). Comparison of indices of traumatic brain injury severity: Glasgow Coma Scale, length of coma and post-traumatic amnesia. *Journal of Neurology, Neurosurgery, and Psychiatry*, 79(6), 678-85. doi:10.1136/jnnp.2006.111187
- SPSS inc. (2013). *IBM SPSS Statistics for Windows*. Armonk, NY: IBM Corporation.
- Thaler, N. S., Bello, D. T., Randall, C., Goldstein, G., Mayfield, J., & Allen, D. N. (2010). IQ profiles are associated with differences in behavioral functioning following pediatric traumatic brain injury. *Archives of Clinical Neuropsychology*, 25(8), 781-90. doi:10.1093/arclin/acq073
- Tsao, J. (2010). *Traumatic Brain Injury: A Clinician's Guide to Diagnosis, Management, and Rehabilitation*. Springer Science & Business Media.
- Wechsler, D. (1955). *Manual for the Wechsler Adult Intelligence Scale*. New York Psychological Corporation.
- Wechsler, D. (1967). *Wechsler preschool and primary scale of intelligence*. Psychological Corporation New York.
- Wechsler, D. (1974). *WISC-R Manual. Wechsler Intelligence Scale for Children-Revised Manual*. Psychological Corporation, New York.
- Wechsler, D. (1981). *WAIS-R manual: Wechsler adult intelligence scale-revised*. Psychological Corporation.
- Wechsler, D. (1989). *WPPSI-R, manual: Wechsler preschool and primary scale of intelligence, revised*. Psychological Corporation New York, NY.
- Wechsler, D. (1991). *WISC-III: Wechsler intelligence scale for children*. Psychological Corporation San Antonio, TX.
- Wechsler, D. (1997). *WAIS-III: Wechsler adult intelligence scale*. Psychological Corporation San Antonio, TX.
- Wechsler, D. (2003). *Wechsler intelligence scale for children--Fourth Edition (WISC-IV)*. San Antonio, TX: The Psychological Corporation.
- Wechsler, D. (2008). *Wechsler Adult Intelligence Scale-Fourth Edition (WAIS-IV)*. San Antonio, TX: NCS Pearson.
- Wells, G. A., Shea, B., O'Connell, D., Peterson, J., Welch, V., Losos, M., & Tugwell, P. (2012). The Newcastle-Ottawa Scale (NOS) for assessing the quality if nonrandomized studies in meta-analyses. (Available from: [http://www.ohri.ca/programs/clinical\\_epidemiology/oxford.asp](http://www.ohri.ca/programs/clinical_epidemiology/oxford.asp)), 2012. doi:10.2307/632432
- Winogron, H. W., Knights, R. M., & Bawden, H. N. (1984). Neuropsychological deficits following head injury in children. *Journal of Clinical Neuropsychology*, 6(3), 269-286.
- Yasuda, S., Wehman, P., Targett, P., Cifu, D., & West, M. (2001). Return to Work for Persons with Traumatic Brain Injury. *American Journal of Physical Medicine & Rehabilitation*, 80, 852-864.

SUPPORTING INFORMATION

Table 1. Overview of studies included for meta-analysis

| Study Methodology          |          |     |          |        |         |           |        |                  |             | Injury Characteristics |             |           |            |         | Demographics |         |                |           |               |      |
|----------------------------|----------|-----|----------|--------|---------|-----------|--------|------------------|-------------|------------------------|-------------|-----------|------------|---------|--------------|---------|----------------|-----------|---------------|------|
| Ref                        | Severity | n   | IQ Index | Design | Control | Subsample | Injury | Wechsler Scale   | GCS measure | LOC measure            | PTA measure | NOS (1-9) | GCS (15-3) | LOC (d) | PTA (d)      | TPI (y) | Age Injury (y) | Males (%) | Education (y) |      |
|                            |          |     |          |        |         |           |        |                  |             |                        |             |           |            |         |              |         |                |           |               |      |
| Subacute phase of recovery |          |     |          |        |         |           |        |                  |             |                        |             |           |            |         |              |         |                |           |               |      |
| 1                          | Mild     | 19  | FPV      | L      | C       | -         | TBI    | WISC-III/WPPSI-R | Admission   | -                      | -           | 7         | 14.1       | -       | -            | 0.5     | 5.2            | 4.7       | 62            | -    |
| 2                          | Mild     | 29  | PV       | L      | OC      | -         | HI     | WISC             | -           | -                      | Retro       | 7         | 14.0       | 0.0     | 0.5          | 0.1     | 9.6            | 9.6       | 76            | -    |
| 3                          | Mild     | 44  | PV       | CS     | C       | -         | CHI    | WISC-R           | -           | -                      | -           | 5         | 14.6       | -       | -            | 0.3     | 10.1           | 9.8       | 52            | -    |
| 4                          | Mild     | 32  | FPV      | L      | -       | -         | TBI    | WISC-R           | Admission   | Days to GCS = 15       | Pro (TPP)   | 2         | 14.0       | 0.0     | 0.5          | 0.5     | 10.8           | 10.8      | 67            | -    |
| 5                          | Mild     | 31  | PV       | CS     | -       | -         | TBI    | WISC-R           | Admission   | Obeys commands         | TPP         | 5         | 14.6       | 0.0     | 0.4          | 0.3     | 11.7           | 11.4      | 74            | -    |
| 6                          | Mild     | 13  | F        | CS     | C       | -         | TBI    | WASI             | -           | -                      | -           | 5         | 14.0       | 0.0     | 0.5          | 0.1     | 13.3           | 13.3      | 62            | -    |
| 7                          | Mild     | 19  | FPV      | CS     | C       | -         | CHI    | WAIS-R/WISC-R    | Admission   | -                      | -           | 6         | 14.0       | 0.0     | 0.5          | 0.1     | 15.3           | 15.2      | 53            | 9.6  |
| Mild                       |          |     |          |        |         |           |        |                  |             | Child Sample           | Mean        | 5.3       | 14.2       | 0.0     | 0.5          | 0.3     | 10.9           | 10.7      | 63.7          | 9.6  |
|                            |          |     |          |        |         |           |        |                  |             |                        | SD          | 1.7       | 0.3        | 0.0     | 0.0          | 0.2     | 3.2            | 3.3       | 9.4           | .    |
|                            |          |     |          |        |         |           |        |                  |             |                        |             |           |            |         |              |         |                |           |               |      |
| 8                          | Mild     | 15  | F        | L      | C       | Women     | HI     | WAIS-R           | -           | -                      | -           | 4         | 14.0       | 0.0     | 0.5          | 0       | 24.2           | 24.2      | 0             | 7.9  |
| 8                          | Mild     | 15  | F        | L      | C       | Men       | HI     | WAIS-R           | -           | -                      | -           | 4         | 14.0       | 0.0     | 0.5          | 0       | 24.7           | 24.7      | 100           | 8.6  |
| 9                          | Mild     | 70  | FPV      | CS     | -       | -         | CHI    | WAIS/WISC-R      | Admission   | Retro (report)         | -           | 4         | -          | 0.0     | -            | 0.3     | 22.6           | 25.3      | 11.5          | -    |
| 10                         | Mild     | 157 | P        | L      | OC      | GCS       | TBI    | WAIS             | Admission   | -                      | Retro       | 7         | 14.4       | -       | -            | 0.1     | 28.5           | 28.4      | 71            | 12.5 |
| 11                         | Mild     | 22  | FPV      | CS     | OC      | -         | TBI    | WAIS-R           | -           | -                      | Retro       | 2         | 14.0       | 0.0     | 0.5          | 0       | 31.2           | 31.2      | 67            | -    |
| 12                         | Mild     | 46  | F        | L      | -       | -         | TBI    | WAIS-R           | -           | -                      | -           | 4         | -          | 0.0     | -            | 0.1     | 33.2           | 33.1      | 78            | 12.1 |
| 13                         | Mild     | 28  | FPV      | CS     | -       | -         | HI     | WISC-R           | -           | -                      | -           | 2         | 14.0       | 0.0     | 0.5          | 0.3     | 35.5           | 35.2      | 78            | -    |
| Mild                       |          |     |          |        |         |           |        |                  |             | Adult Sample           | Mean        | 3.9       | 14.1       | 0.0     | 0.5          | 0.1     | 28.6           | 28.9      | 65.7          | 10.5 |
|                            |          |     |          |        |         |           |        |                  |             |                        | SD          | 1.7       | 0.2        | 0.0     | 0.0          | 0.1     | 4.9            | 4.4       | 34.1          | 2.1  |
|                            |          |     |          |        |         |           |        |                  |             |                        |             |           |            |         |              |         |                |           |               |      |
| Total                      |          | 540 |          |        |         |           |        |                  |             |                        | Mean        | 4.6       | 14.1       | 0.0     | 0.5          | 0.2     | 19.7           | 19.8      | 64.6          | 10.4 |
|                            |          |     |          |        |         |           |        |                  |             |                        | SD          | 1.8       | 0.2        | 0.0     | 0.0          | 0.2     | 10.0           | 10.1      | 23.0          | 1.9  |

| Study Methodology |          |     |          |        |         |           |        |                  |              |                  | Injury Characteristics |           |                |         |         | Demographics |         |                |           |               |
|-------------------|----------|-----|----------|--------|---------|-----------|--------|------------------|--------------|------------------|------------------------|-----------|----------------|---------|---------|--------------|---------|----------------|-----------|---------------|
| Ref               | Severity | n   | IQ Index | Design | Control | Subsample | Injury | Wechsler Scale   | GCS measure  | LOC measure      | PTA measure            | NOS (1-9) | GCS (15-3) (d) | LOC (d) | PTA (d) | TPI (y)      | Age (y) | Age Injury (y) | Males (%) | Education (y) |
| 1                 | Mod      | 46  | FPV L    | C      | -       | -         | TBI    | WISC-III/WPPSI-R | Admission    | -                | -                      | 7         | 10.3           | -       | -       | 0.2          | 4.7     | 4.5            | 62        | -             |
| 4                 | Mod      | 18  | FPV L    | -      | -       | -         | TBI    | WISC-R           | Admission    | Days to GCS = 15 | Pro (TPP)              | 2         | 11.2           | -       | -       | 0.5          | 10.8    | 10.8           | 67        | -             |
| 5                 | Mod      | 7   | PV CS    | -      | -       | -         | TBI    | WISC-R           | Admission    | Obey commands    | Pro (TPP)              | 5         | 10.3           | -       | 2.6     | 0.2          | 11.1    | 10.9           | 57        | -             |
| 14                | Mod      | 30  | FPV CS   | OC     | -       | -         | CHI    | WISC-III         | -            | -                | COAT                   | 4         | -              | -       | 4       | 0            | 10.9    | 10.9           | 67        | -             |
| Mod               |          |     |          |        |         |           |        |                  | Child sample |                  | Mean                   | 4.5       | 10.6           | -       | 3.3     | 0.2          | 9.4     | 9.3            | 63.3      |               |
|                   |          |     |          |        |         |           |        |                  |              |                  | SD                     | 2.1       | 0.5            | -       | 1.0     | 0.2          | 3.1     | 3.2            | 4.9       |               |
|                   |          |     |          |        |         |           |        |                  |              |                  |                        |           |                |         |         |              |         |                |           |               |
| 15                | Mod      | 17  | FPV CS   | -      | -       | -         | CHI    | WAIS-R           | -            | -                | Retro                  | 3         | 9.4            | -       | 15.2    | 0.5          | 42.9    | 42.5           | 53        | 12.1          |
| 16                | Mod      | 4   | FPV CS   | -      | MMI     | -         | BI     | WAIS-R           | Admission    | -                | GOAT                   | 5         | 11.0           | -       | 5       | 0            | 47.0    | 47.0           | 75        | -             |
| Mod               |          |     |          |        |         |           |        |                  |              | Adult Sample     |                        | 4.3       | 10.2           | -       | 6.7     | 0.2          | 21.2    | 21.1           | 63.5      | 12.1          |
|                   |          |     |          |        |         |           |        |                  |              |                  | SD                     | 1.8       | 1.1            | -       | 5.8     | 0.2          | 18.6    | 18.5           | 7.9       | 0.2           |
|                   |          |     |          |        |         |           |        |                  |              |                  |                        |           |                |         |         |              |         |                |           |               |
| Total             |          | 122 |          |        |         |           |        |                  |              |                  | Mean                   | 4.6       | 10.4           | 5.4     | 6.7     | 0.2          | 22.6    | 20.2           | 66.0      | 12.0          |
|                   |          |     |          |        |         |           |        |                  |              |                  | SD                     | 1.8       | 0.7            | 1.5     | 5.8     | 0.2          | 17.9    | 16.1           | 8.1       | 0.2           |
|                   |          |     |          |        |         |           |        |                  |              |                  |                        |           |                |         |         |              |         |                |           |               |
| 1                 | Severe   | 31  | FPV L    | C      | -       | -         | TBI    | WISC-III/WPPSI-R | Admission    | -                | -                      | 7         | 5.1            | -       | -       | 0.3          | 4.9     | 4.7            | 62        | -             |
| 17                | Severe   | 16  | FPV L    | -      | Early   | -         | HI     | WISC-R/WPPSI-R   | Lowest 24h   | Days to GCS = 8  | Orientation            | 4         | 5.3            | 9.2     | 14.6    | 0.4          | 6       | 5.7            | 59        | -             |
| 18                | Severe   | 42  | FPV CS   | -      | -       | -         | TBI    | WISC-R           | Admission    | -                | -                      | 4         | -              | 4.5     | -       | 0.3          | 8.4     | 8.2            | 74        | -             |
| 3                 | Severe   | 68  | PV CS    | C      | -       | -         | CHI    | WISC-R           | -            | -                | -                      | 5         | 5.8            | -       | -       | 0.4          | 9.9     | 9.5            | 59        | -             |
| 17                | Severe   | 20  | FPV L    | -      | Late    | -         | HI     | WISC-R/WPPSI-R   | Lowest 24h   | Days to GCS = 8  | Orientation            | 4         | 5.2            | 7.8     | 18      | 0.4          | 11.2    | 10.8           | 59        | -             |
| 4                 | Severe   | 26  | FPV L    | -      | -       | -         | TBI    | WISC-R           | Admission    | Days to GCS = 15 | Pro (TPP)              | 2         | 5.5            | 25.2    | 54.2    | 0.5          | 10.8    | 10.8           | 67        | -             |
| 19                | Severe   | 50  | F CS     | OC     | -       | -         | TBI    | WASI             | Lowest       | -                | -                      | 7         | -              | 6.4     | -       | 0.2          | 11.3    | 11.1           | 74        | -             |
| 5                 | Severe   | 13  | PV CS    | -      | -       | -         | TBI    | WISC-R           | Admission    | Obey commands    | TPP                    | 5         | 5.0            | 3.3     | 20.8    | 0.3          | 11.6    | 11.4           | 69        | -             |
| 20                | Severe   | 17  | FPV L    | -      | -       | -         | TBI    | WISC             | -            | -                | -                      | 4         | -              | 17.6    | -       | 0.2          | 12.0    | 11.8           | 39        | -             |
| 21                | Severe   | 30  | F CS     | -      | -       | -         | CHI    | WISC-R           | -            | -                | -                      | 3         | -              | 11.7    | -       | 0.2          | 11.9    | 11.8           | 53        | -             |
| 22                | Severe   | 77  | FPV CS   | -      | -       | -         | TBI    | WISC-III         | -            | -                | -                      | 5         | 6.0            | 24.1    | -       | 0.4          | 12.3    | 11.8           | 74        | -             |



| Study Methodology              |          |    |          |        |         |           |        |                |             | Injury Characteristics |             |           |            |         |         | Demographics |         |                |           |               |      |
|--------------------------------|----------|----|----------|--------|---------|-----------|--------|----------------|-------------|------------------------|-------------|-----------|------------|---------|---------|--------------|---------|----------------|-----------|---------------|------|
| Ref                            | Severity | n  | IQ Index | Design | Control | Subsample | Injury | Wechsler Scale | GCS measure | LOC measure            | PTA measure | NOS (1-9) | GCS (15-3) | LOC (d) | PTA (d) | TPI (y)      | Age (y) | Age Injury (y) | Males (%) | Education (y) |      |
| 23                             | Severe   | 20 | P V      | L      | -       | -         | TBI    | WISC           | Admission   | -                      | -           | 4         | 5.0        | 16      | 69      | 0.5          | 12.2    | 12.2           | 65        | -             |      |
|                                |          |    |          |        |         |           |        |                |             | Child Sample           |             | Mean      | 4.5        | 5.4     | 12.6    | 35.3         | 0.3     | 10.2           | 10.0      | 62.8          | -    |
|                                |          |    |          |        |         |           |        |                |             | SD                     |             | SD        | 1.4        | 0.4     | 7.8     | 24.7         | 0.1     | 2.5            | 2.5       | 10.2          | -    |
| 24                             | Severe   | 16 | P V      | L      | -       | Not RE    | TBI    | WAIS-R         | Admission   | -                      | -           | 4         | -          | -       | 107.8   | 0.5          | 23.3    | 22.8           | 62        | 11.8          |      |
| 25                             | Severe   | 52 | F        | L      | -       | -         | TBI    | WAIS-R         | -           | -                      | -           | 4         | -          | 4.3     | -       | 0.2          | 42.2    | 24.1           | 73        | 11.8          |      |
| 26                             | Severe   | 30 | F        | CS     | -       | -         | HI     | WAIS           | -           | -                      | -           | 5         | -          | 54.8    | -       | 0.5          | 24.7    | 24.2           | 70        | 13.1          |      |
| 24                             | Severe   | 19 | P V      | L      | -       | RE        | TBI    | WAIS-R         | Admission   | -                      | -           | 4         | -          | -       | 28      | 0.3          | 24.3    | 24.5           | 79        | 13.1          |      |
| 12                             | Severe   | 27 | F        | L      | -       | -         | TBI    | WAIS-R         | -           | -                      | -           | 4         | -          | 20.1    | -       | 0.2          | 24.8    | 24.6           | 67        | 11.9          |      |
| 27                             | Severe   | 8  | FPV      | CS     | -       | -         | TBI    | WAIS-III       | Admission   | -                      | -           | 3         | 7.5        | -       | -       | 0.2          | 26.0    | 25.8           | 88        | -             |      |
| 28                             | Severe   | 23 | FPV      | CS     | C       | -         | CHI    | WAIS-R         | Lowest      | -                      | Westmead    | 6         | 9.0        | -       | 28.5    | 0            | 26.2    | 26.1           | 81        | 10.0          |      |
| 29                             | Severe   | 39 | FPV      | CS     | -       | -         | CHI    | WAIS-R         | -           | -                      | OGMS        | 3         | -          | -       | 66.8    | <0.5         | 27.6    | 27.6           | -         | 11.7          |      |
| 30                             | Severe   | 10 | FPV      | CS     | C       | -         | HI     | WAIS           | -           | -                      | Aware 24h   | 6         | -          | -       | 22      | 0.5          | 29.0    | 28.5           | 92        | -             |      |
| 31                             | Severe   | 15 | FPV      | CS     | -       | -         | TBI    | WAIS-R         | Lowest 24h  | Obey commands          | -           | 4         | -          | 3.3     | -       | 0.4          | 33.4    | 33.0           | 64        | 12.0          |      |
| 16                             | Severe   | 6  | FPV      | CS     | -       | SMI       | BI     | WAIS-R         | Admission   | -                      | GOAT        | 5         | 9.0        | -       | 10      | 0            | 35.1    | 35             | 83        | -             |      |
| 16                             | Severe   | 20 | FPV      | CS     | -       | SMI       | BI     | WAIS-R         | Admission   | -                      | GOAT        | 5         | 9.0        | -       | 12      | 0            | 38.1    | 38.0           | 70        | -             |      |
| 16                             | Severe   | 7  | FPV      | CS     | -       | MMI       | BI     | WAIS-R         | Admission   | -                      | GOAT        | 5         | 8.0        | -       | 9       | 0            | 41.1    | 41.0           | 71        | -             |      |
| 32                             | Severe   | 50 | FPV      | CS     | -       | -         | CHI    | WAIS-R         | -           | -                      | NA          | 3         | 10.6       | -       | 29.7    | 0.1          | 41.5    | 41.5           | 74        | 11.6          |      |
|                                |          |    |          |        |         |           |        |                |             | Adult Sample           |             | Mean      | 4.4        | 8.9     | 20.6    | 34.9         | 0.2     | 31.2           | 29.8      | 74.9          | 11.9 |
|                                |          |    |          |        |         |           |        |                |             | SD                     |             | SD        | 1.0        | 1.1     | 24.0    | 32.5         | 0.2     | 7.1            | 6.6       | 9.1           | 0.9  |
|                                |          |    |          |        |         |           |        |                |             |                        |             |           |            |         |         |              |         |                |           |               |      |
| Total 732                      |          |    |          |        |         |           |        |                |             | Mean                   |             | 4.4       | 7.1        | 14.9    | 34.7    | 0.3          | 22.3    | 21.4           | 69.3      | 11.9          |      |
|                                |          |    |          |        |         |           |        |                |             | SD                     |             | 1.2       | 1.9        | 1.8     | 28.9    | 0.2          | 12.0    | 11.3           | 11.3      | 0.9           |      |
|                                |          |    |          |        |         |           |        |                |             |                        |             |           |            |         |         |              |         |                |           |               |      |
| Total TBI severity groups 1394 |          |    |          |        |         |           |        |                |             | Mean                   |             | 4.5       | 10.3       | 8.3     | 19.3    | 0.3          | 20.9    | 20.4           | 67.0      | 11.3          |      |
|                                |          |    |          |        |         |           |        |                |             | SD                     |             | 1.5       | 3.6        | 11.8    | 12.1    | 15.2         | 1.4     | 1.5            | 15.2      | 1.5           |      |



| Study Methodology         |          |     |          |        |         |             |              |                  |             | Injury Characteristics |             |           |                |         | Demographics |         |                |           |               |      |
|---------------------------|----------|-----|----------|--------|---------|-------------|--------------|------------------|-------------|------------------------|-------------|-----------|----------------|---------|--------------|---------|----------------|-----------|---------------|------|
| Ref                       | Severity | n   | IQ Index | Design | Control | Subsample   | Injury       | Wechsler Scale   | GCS measure | LOC measure            | PTA measure | NOS (1-9) | GCS (15-3) (d) | LOC (d) | PTA (d)      | TPI (y) | Age Injury (y) | Males (%) | Education (y) |      |
|                           |          |     |          |        |         |             |              |                  |             |                        |             |           |                |         |              |         |                |           |               |      |
| Chronic phase of recovery |          |     |          |        |         |             |              |                  |             |                        |             |           |                |         |              |         |                |           |               |      |
| 1                         | Mild     | 19  | FPV      | L      | C       | -           | TBI          | WISC-III/WPPSI-R | Admission   | -                      | -           | 7         | 14.1           | -       | -            | 1.5     | 6.2            | 4.7       | 62            | -    |
| 33                        | Mild     | 15  | F        | CS     | C       | -           | TBI          | WISC-III         | Admission   | -                      | -           | 7         | 15.0           | 0.0     | 0.0          | 1       | 8.3            | 7.3       | 33            | -    |
| 34                        | Mild     | 67  | PV       | CS     | -       | -           | HI           | WISC-R           | Stable GCS  | -                      | -           | 4         | 14.7           | -       | -            | 2.4     | 10.0           | 7.6       | -             | -    |
| 2                         | Mild     | 29  | PV       | L      | OC      | -           | HI           | WISC             | -           | Retro                  | -           | 7         | 14.0           | 0.0     | 0.5          | 1       | 10.6           | 9.6       | 76            | -    |
| 35                        | Mild     | 24  | FPV      | CS     | OC      | -           | TBI          | WISC-R           | Lowest      | Obey commands          | -           | 5         | 14.5           | 0.0     | -            | 2.4     | 11.2           | 8.8       | 75            | -    |
| 4                         | Mild     | 32  | FPV      | L      | -       | -           | TBI          | WISC-R           | Admission   | Days to GCS = 15       | Pro (TPP)   | 2         | 14.0           | 0.0     | 0.5          | 0.8     | 11.6           | 10.8      | 67            | -    |
| 36                        | Mild     | 31  | FPV      | CS     | C       | -           | TBI          | WISC-III         | -           | -                      | -           | 6         | 14.0           | 0.0     | 0.5          | 2       | 11.7           | 8.9       | 60            | -    |
| 37                        | Mild     | 7   | V&P      | CS     | -       | Adolescents | CHI          | WISC/WAIS        | Admission   | -                      | -           | 3         | -              | 0.03    | -            | 1.3     | 17.8           | 16.5      | -             | -    |
| Mild                      |          |     |          |        |         |             | Child Sample |                  |             |                        | Mean        | 5.1       | 14.3           | 0.0     | 0.4          | 1.6     | 10.9           | 9.3       | 62.2          | -    |
|                           |          |     |          |        |         |             | SD           |                  |             |                        | SD          | 2.0       | 0.4            | 0.0     | 0.3          | 0.6     | 3.4            | 3.4       | 15.7          | -    |
| 8                         | Mild     | 15  | F        | L      | C       | Women       | HI           | WAIS-R           | -           | -                      | -           | 4         | 14.0           | 0.0     | 0.5          | 0.5     | 24.2           | 24.2      | 0             | 7.9  |
| 8                         | Mild     | 15  | F        | L      | C       | Men         | HI           | WAIS-R           | -           | -                      | -           | 4         | 14.0           | 0.0     | 0.5          | 0.5     | 24.7           | 24.7      | 100           | 8.6  |
| 10                        | Mild     | 157 | P        | L      | OC      | GCS         | TBI          | WAIS             | Admission   | -                      | Retro       | 7         | 14.4           | -       | -            | 1       | 29.4           | 28.4      | 71            | 12.5 |
| 25                        | Mild     | 46  | F        | L      | -       | -           | TBI          | WAIS-R           | -           | -                      | -           | 4         | -              | 0.0     | -            | 1.03    | 34.1           | 33.1      | 78            | 12.1 |
| 38                        | Mild     | 22  | FPV      | CS     | C       | -           | CHI          | WAIS-R           | -           | -                      | -           | 5         | 14.0           | 0.0     | 0.5          | 1.1     | 34.3           | 33.3      | 64            | 14.6 |
| 39                        | Mild     | 41  | FPV      | CS     | -       | -           | CHI          | WAIS-R           | -           | -                      | -           | 2         | 14.0           | 0.0     | 0.5          | 1.8     | 34.5           | 32.6      | 12.8          | 12.8 |
| 40                        | Mild     | 42  | V        | CS     | -       | -           | TBI          | WAIS-III         | -           | -                      | -           | 3         | 14.0           | 0.0     | 0.5          | 8.8     | 35.6           | 26.8      | 45            | 12.9 |
| 41                        | Mild     | 23  | FPV      | CS     | C       | -           | TBI          | WAIS-III         | -           | -                      | -           | 5         | -              | 0.0     | 0.0          | 1.18    | 35.7           | 34.6      | 52            | 12.9 |
| 42                        | Mild     | 26  | FPV      | CS     | -       | -           | TBI          | WAIS-III         | -           | -                      | -           | 3         | 14.0           | 0.0     | 0.5          | 1.12    | 38.6           | 37.5      | 73            | 12.0 |
| 12                        | Mild     | 22  | F        | CS     | -       | -           | TBI          | WAIS-III         | Admission   | -                      | -           | 4         | 14.6           | -       | -            | -       | 43.2           | -         | 59            | 13.1 |
| 43                        | Mild     | 34  | F,P,V    | CS     | -       | -           | HI           | WAIS-R/-III      | -           | -                      | -           | 4         | -              | 0.0     | -            | ≥1      | 45.4           | -         | 73            | 12.0 |
| 44                        | Mild     | 29  | FPV      | CS     | -       | Women       | TBI          | K-WAIS           | -           | -                      | -           | 4         | 14.0           | 0.0     | 0.5          | 1.9     | 45.7           | 43.8      | 0             | 9.7  |
| 44                        | Mild     | 95  | FPV      | CS     | -       | Men         | TBI          | K-WAIS           | -           | -                      | -           | 4         | 14.0           | 0.0     | 0.5          | 1.9     | 45.7           | 43.8      | 100           | 9.7  |
| Mild                      |          |     |          |        |         |             | Adult Sample |                  |             |                        | Mean        | 4.1       | 14.1           | 0.0     | 0.4          | 1.9     | 37.5           | 33.0      | 61.5          | 11.7 |
|                           |          |     |          |        |         |             | SD           |                  |             |                        | SD          | 1.1       | 0.2            | 0.0     | 0.2          | 2.3     | 7.6            | 6.8       | 30.1          | 1.8  |

| Study Methodology |          |     |          |        |         |           |        |                   |             | Injury Characteristics |             |           |            |         | Demographics |         |         |                |           |               |
|-------------------|----------|-----|----------|--------|---------|-----------|--------|-------------------|-------------|------------------------|-------------|-----------|------------|---------|--------------|---------|---------|----------------|-----------|---------------|
| Ref               | Severity | n   | IQ Index | Design | Control | Subsample | Injury | Wechsler Scale    | GCS measure | LOC measure            | PTA measure | NOS (1-9) | GCS (15-3) | LOC (d) | PTA (d)      | TPI (y) | Age (y) | Age Injury (y) | Males (%) | Education (y) |
|                   |          |     |          |        |         |           |        |                   |             |                        |             |           |            |         |              |         |         |                |           |               |
| Total             |          | 859 |          |        |         |           |        |                   |             |                        | Mean        | 4.5       | 14.2       | 0.0     | 0.4          | 1.7     | 26.6    | 23.0           | 60.4      | 11.6          |
|                   |          |     |          |        |         |           |        |                   |             |                        | SD          | 1.6       | 0.3        | 0.0     | 0.2          | 1.8     | 14.0    | 13.2           | 27.4      | 2.0           |
| <sup>1</sup>      | Mod      | 46  | FPV      | L      | C       | -         | TBI    | WISC-III/WPPSI-R  | Admission   | -                      | -           | 7         | 10.3       | -       | -            | 1.2     | 5.7     | 4.5            | 62        | -             |
| <sup>45</sup>     | Mod      | 43  | PV       | CS     | -       | -         | HI     | WISC-III          | -           | -                      | -           | 3         | 10.6       | -       | -            | 4       | 10.9    | 7              | 61        | -             |
| <sup>34</sup>     | Mod      | 14  | PV       | CS     | -       | -         | HI     | WISC-R            | Stable GCS  | -                      | -           | 4         | 9.9        | -       | -            | 2.4     | 10      | 7.6            | -         | -             |
| <sup>46</sup>     | Mod      | 19  | F        | CS     | -       | EI        | TBI    | WISC-IV           | Admission   | -                      | -           | 3         | 10.6       | -       | -            | 3.2     | 12.6    | 9.4            | 68        | -             |
| <sup>37</sup>     | Mod      | 9   | V&P      | CS     | -       | Children  | CHI    | WISC/WAIS         | Admission   | -                      | -           | 3         | -          | 0.5     | -            | 2.9     | 11.7    | 8.8            | -         | -             |
| <sup>4</sup>      | Mod      | 18  | FPV      | L      | -       | -         | TBI    | WISC-R            | Admission   | Days to GCS = 15       | Pro (TPP)   | 2         | 11.2       | -       | -            | 0.8     | 11.6    | 10.8           | 67        | -             |
| <sup>47</sup>     | Mod      | 17  | FPV      | CS     | C       | -         | HI     | WISC-R/-III       | Admission   | Days to GCS = 9        | -           | 5         | 10.6       | -       | -            | 3.2     | 14.9    | 11.9           | 47        | -             |
| <sup>48</sup>     | Mod      | 20  | F        | CS     | -       | TOPS      | TBI    | WASI              | Lowest      | -                      | -           | 4         | 6.5        | -       | -            | 0.7     | 14      | 13.5           | 38        | -             |
| <sup>48</sup>     | Mod      | 21  | F        | CS     | -       | IRC       | TBI    | WASI              | Lowest      | -                      | -           | 4         | 10.5       | -       | -            | 0.9     | 14.5    | 13.7           | 58        | -             |
| <sup>49</sup>     | Mod      | 14  | F        | CS     | C       | -         | TBI    | WISC-IV/WAIS-III  | Lowest      | -                      | -           | 5         | 12.3       | -       | -            | 0.7     | 15.1    | 14.4           | 36        | -             |
| Mod               |          |     |          |        |         |           |        |                   |             | Child Sample           | Mean        | 4.0       | 10.3       | 0.5     | -            | 2.0     | 12.1    | 10.2           | 54.6      |               |
|                   |          |     |          |        |         |           |        |                   |             |                        | SD          | 1.4       | 1.6        | -       | -            | 1.3     | 2.9     | 3.3            | 12.6      |               |
| <sup>50</sup>     | Mod      | 115 | F        | CS     | -       | -         | HI     | WPPSI/WISC-R/WAIS | 6h GCS      | -                      | -           | 4         | 9.6        | -       | -            | 1.8     | 21.6    | 19.8           | 75        | -             |
| <sup>51</sup>     | Mod      | 18  | FPV      | CS     | -       | -         | TBI    | WAIS-III          | -           | -                      | -           | 3         | 11.9       | -       | -            | 2       | 28.1    | 26.1           | 61        | -             |
| <sup>52</sup>     | Mod      | 30  | FPV      | CS     | -       | Control   | TBI    | WAIS-II/WAIS-R    | -           | -                      | -           | 2         | -          | 0.09    | 3.3          | 2       | 33.6    | 31.3           | 70        | 12.52         |
| <sup>53</sup>     | Mod      | 107 | FPV      | CS     | -       | -         | TBI    | Wechsler          | -           | -                      | -           | 2         | -          | 0.13    | -            | 0.8     | 34.1    | 33.3           | 79        | 11.6          |
| Mod               |          |     |          |        |         |           |        |                   |             | Adult Sample           | Mean        | 2.8       | 10.8       | 0.1     | 3.3          | 1.7     | 29.4    | 27.6           | 71.3      | 12.1          |
|                   |          |     |          |        |         |           |        |                   |             |                        | SD          | 1.0       | 1.6        | 0.0     | -            | 0.6     | 5.8     | 6.0            | 7.8       | 0.7           |
| Total             |          | 491 |          |        |         |           |        |                   |             |                        | Mean        | 3.5       | 10.1       | 0.2     | 3.3          | 2.1     | 17.4    | 15.3           | 62.8      | 12.1          |
|                   |          |     |          |        |         |           |        |                   |             |                        | SD          | 1.4       | 1.5        | 0.2     | -            | 1.1     | 9.6     | 9.9            | 12.4      | 0.7           |



| Study Methodology |          |     |          |        |         |              |        |                  |             | Injury Characteristics |             |             |         |      |             | Demographics |           |            |      |        |         |         |         |         |                |           |               |
|-------------------|----------|-----|----------|--------|---------|--------------|--------|------------------|-------------|------------------------|-------------|-------------|---------|------|-------------|--------------|-----------|------------|------|--------|---------|---------|---------|---------|----------------|-----------|---------------|
| Ref               | Severity | n   | IQ Index | Design | Control | Subsample    | Injury | Wechsler Scale   | GCS measure |                        |             | LOC measure |         |      | PTA measure |              | NOS (1-9) | GCS (15-3) |      |        | LOC (d) | PTA (d) | TPI (y) | Age (y) | Age Injury (y) | Males (%) | Education (y) |
|                   |          |     |          |        |         |              |        |                  | GCS         | measure                |             | LOC         | measure |      | PTA         | measure      |           |            | GCS  | (15-3) |         |         |         |         |                |           |               |
| 1                 | Severe   | 31  | FPV      | L      | C       | -            | TBI    | WISC-III/WPPSI-R | Admission   | -                      | -           | 7           | 5.1     | -    | -           | 1.3          | 5.9       | 4.7        | 61   | -      |         |         |         |         |                |           |               |
| 17                | Severe   | 16  | FPV      | L      | -       | Early injury | HI     | WISC-R/WPPSI-R   | Lowest 24h  | Days to GCS = 8        | Orientation | 4           | 5.3     | 9.2  | 14.6        | 2.0          | 7.7       | 5.7        | 58   | -      |         |         |         |         |                |           |               |
| 37                | Severe   | 15  | PV       | CS     | -       | Children     | CHI    | WAIS             | Admission   | -                      | -           | 3           | -       | 21   | -           | 1.1          | 8.6       | 7.5        | -    | -      |         |         |         |         |                |           |               |
| 54                | Severe   | 14  | F        | CS     | C       | Children     | TBI    | WASI             | Admission   | -                      | -           | 6           | 8.3     | -    | -           | 3.9          | 9.1       | 5.2        | 57   | -      |         |         |         |         |                |           |               |
| 34                | Severe   | 5   | PV       | CS     | -       | -            | HI     | WISC-R           | Stable GCS  | -                      | -           | 4           | 5.4     | -    | -           | 2.4          | 10.0      | 7.6        | -    | -      |         |         |         |         |                |           |               |
| 35                | Severe   | 23  | FPV      | CS     | OC      | -            | TBI    | WISC-R           | Lowest      | ObeY commands          | -           | 5           | 5.3     | 17.9 | -           | 2.0          | 10.7      | 8.71       | 75   | -      |         |         |         |         |                |           |               |
| 55                | Severe   | 23  | FPV      | CS     | -       | -            | TBI    | WISC             | Admission   | -                      | -           | 3           | 5.3     | -    | -           | 4.1          | 10.7      | 6.6        | 57   | 4.2    |         |         |         |         |                |           |               |
| 56                | Severe   | 76  | V        | CS     | -       | -            | TBI    | WISC-III         | Admission   | -                      | -           | 4           | 5.9     | -    | -           | 1.0          | 11.1      | 10.1       | 61   | -      |         |         |         |         |                |           |               |
| 57                | Severe   | 20  | FPV      | CS     | C       | -            | TBI    | WISC-R           | -           | -                      | -           | 4           | -       | 15.8 | 50.5        | 1.7          | 11.1      | 9.6        | 70   | -      |         |         |         |         |                |           |               |
| 58                | Severe   | 36  | F        | CS     | C       | -            | TBI    | WISC-III         | Lowest      | -                      | Westmead    | 6           | 6.7     | 16.8 | 26.5        | 4.3          | 11.3      | 6.6        | 67   | -      |         |         |         |         |                |           |               |
| 4                 | Severe   | 26  | FPV      | L      | -       | -            | TBI    | WISC-R           | Admission   | Days to GCS = 15       | Pro (TPP)   | 2           | 5.5     | 25.2 | 54.2        | 0.8          | 11.6      | 10.8       | 67   | -      |         |         |         |         |                |           |               |
| 59                | Severe   | 17  | FPV      | SC     | SC      | -            | HI     | WISC-R           | Admission   | -                      | -           | 5           | 8.8     | 6    | -           | 3.4          | 11.9      | 8.3        | 67   | -      |         |         |         |         |                |           |               |
| 46                | Severe   | 9   | F        | CS     | -       | PI           | TBI    | WISC-IV          | Admission   | -                      | -           | 3           | 6.7     | -    | -           | 3.4          | 12.3      | 9.17       | 78   | -      |         |         |         |         |                |           |               |
| 17                | Severe   | 20  | FPV      | L      | -       | Late injury  | HI     | WISC-R, WPPSI-R  | Lowest 24h  | Days to GCS = 8        | Orientation | 4           | 5.2     | 7.8  | 18          | 2.0          | 12.8      | 10.8       | 58   | -      |         |         |         |         |                |           |               |
| 60                | Severe   | 151 | F        | CS     | C       | -            | TBI    | WISC-III         | Lowest 24h  | -                      | -           | 7           | 8.0     | -    | -           | 0.7          | 12.9      | 12.2       | 58   | -      |         |         |         |         |                |           |               |
| 61                | Severe   | 40  | FPV      | CS     | -       | TBI baseline | TBI    | WISC-R/WAIS-R    | -           | -                      | -           | 4           | -       | 21.1 | -           | 0.6          | 13.9      | 13.4       | 75   | 6      |         |         |         |         |                |           |               |
| 54                | Severe   | 14  | F        | CS     | C       | Adolescents  | TBI    | WASI             | Admission   | -                      | -           | 6           | 6.5     | -    | -           | 4.9          | 14.3      | 9.4        | 86   | -      |         |         |         |         |                |           |               |
| 47                | Severe   | 16  | FPV      | CS     | C       | -            | HI     | WISC-II/ WISC-R  | Admission   | Days to GCS = 9        | -           | 5           | 6.1     | 10.1 | -           | 5.3          | 15.0      | 9          | 47   | -      |         |         |         |         |                |           |               |
| 61                | Severe   | 25  | FPV      | CS     | -       | Cbaseline    | TBI    | WISC-R/WAIS-R    | -           | -                      | -           | 4           | -       | 20.6 | -           | 0.8          | 15.5      | 13.5       | 64   | 5.7    |         |         |         |         |                |           |               |
| 37                | Severe   | 14  | PV       | CS     | -       | Adolescents  | CHI    | WAIS             | Admission   | -                      | -           | 3           | -       | 28.3 | -           | 0.6          | 17.5      | 16.9       | -    | -      |         |         |         |         |                |           |               |
| 62                | Severe   | 19  | FPV      | CS     | C       | -            | TBI    | WAIS             | Admission   | -                      | -           | 5           | 5.9     | -    | -           | 9.4          | 8.6       | 18.1       | 86   | -      |         |         |         |         |                |           |               |
| 20                | Severe   | 16  | FPV      | L      | -       | -            | TBI    | WISC             | -           | -                      | -           | 4           | -       | 17.6 | -           | 6.8          | 18.8      | 11.8       | 39   | -      |         |         |         |         |                |           |               |
| 23                | Severe   | 20  | PV       | L      | -       | WAIS         | TBI    | WAIS             | Admission   | -                      | -           | 4           | 5.0     | 16   | 69          | 9.7          | 22.3      | 12.2       | 65   | -      |         |         |         |         |                |           |               |
| 24*               | Severe   | 16  | PV       | L      | -       | Not RE       | TBI    | WAIS-R           | Admission   | -                      | -           | 4           | -       | -    | 107.8       | 1.7          | 24.5      | 22.8       | 62   | 11.8   |         |         |         |         |                |           |               |
| Severe            |          |     |          |        |         |              |        |                  |             | Child Sample           | Mean        | 4.4         | 6.2     | 16.7 | 48.7        | 2.8          | 13.0      | 10.1       | 63.6 | 6.9    |         |         |         |         |                |           |               |
|                   |          |     |          |        |         |              |        |                  |             | SD                     | SD          | 1.3         | 1.2     | 6.5  | 33.0        | 2.3          | 4.4       | 4.0        | 10.6 | 3.3    |         |         |         |         |                |           |               |

| Study Methodology |            |     |          |        |         |           |        |                |             | Injury Characteristics |             |           |            |         |         | Demographics |                |           |               |      |
|-------------------|------------|-----|----------|--------|---------|-----------|--------|----------------|-------------|------------------------|-------------|-----------|------------|---------|---------|--------------|----------------|-----------|---------------|------|
| Ref               | Severity   | n   | IQ Index | Design | Control | Subsample | Injury | Wechsler Scale | GCS measure | LOC measure            | PTA measure | NOS (1-9) | GCS (15-3) | LOC (d) | PTA (d) | TPI (y)      | Age Injury (y) | Males (%) | Education (y) |      |
| 63                | Severe 16  | F   | CS       | C      | Young   |           | TBI    | WAIS-III       | Admission   | -                      | -           | 4         | 5.4        | -       | 22.1    | 6.1          | 24.7           | 18.6      | 94            | -    |
| 24                | Severe 19  | PV  | L        | -      | RE      |           | TBI    | WAIS-R         | Admission   | -                      | -           | 4         | -          | -       | 28      | 1.2          | 25.7           | 24.5      | 79            | 13.1 |
| 25                | Severe 27  | F   | L        | -      | -       |           | TBI    | WAIS-R         | -           | -                      | -           | 4         | -          | 20.1    | -       | 1.3          | 25.9           | 24.6      | 67            | 11.9 |
| 23                | Severe 19  | FPV | CS       | -      | RTS     |           | TBI    | WAIS-R         | -           | -                      | -           | 3         | -          | 11.3    | -       | 3.1          | 27.0           | 23.9      | 85            | -    |
| 64*               | Severe 44  | FPV | CS       | -      | -       |           | CHI    | WAIS-III       | -           | -                      | GOAT        | 4         | -          | 45.3    | 110.5   | 0.9          | 27.4           | 26.5      | 71            | 11.7 |
| 65                | Severe 14  | V   | CS       | C      | -       |           | CHI    | WAIS-R         | Lowest      | Retro (records)        | -           | 5         | 5.2        | -       | -       | 4.2          | 27.4           | 23.2      | 71            | 12.9 |
| 25                | Severe 52  | F   | L        | -      | -       |           | TBI    | WAIS-R         | -           | -                      | -           | 4         | -          | 4.3     | -       | 1            | 25.1           | 24.1      | 73            | 11.8 |
| 66                | Severe 63  | PV  | CS       | -      | -       |           | TBI    | WAIS-R         | -           | -                      | -           | 3         | -          | 31.9    | -       | 3.4          | 28.2           | 24.8      | 83            | 12.9 |
| 67                | Severe 66  | F   | CS       | -      | -       |           | TBI    | WAIS-R         | -           | Retro (records)        | -           | 4         | -          | 13.3    | -       | 3.9          | 28.4           | 24.6      | -             | 13.4 |
| 68                | Severe 64  | FPV | CS       | -      | -       |           | CHI    | WAIS-R         | Admission   | -                      | -           | 3         | 8.4        | -       | -       | 1.9          | 29.0           | 27.2      | 52            | 12.7 |
| 69*               | Severe 20  | F   | CS       | C      | -       |           | CHI    | WAIS-R         | -           | -                      | Retro       | 4         | -          | -       | 12.5    | 5.4          | 29.3           | 23.9      | 70            | 14.1 |
| 70                | Severe 55  | F   | CS       | -      | -       |           | CHI    | WAIS-R         | -           | -                      | -           | 2         | -          | -       | 67.5    | 7.4          | 29.3           | 21.9      | 79            | -    |
| 71                | Severe 44  | F   | CS       | -      | -       |           | TBI    | WASI           | Admission   | -                      | Westmead    | 4         | 7.1        | -       | 47.9    | >5.0         | 29.6           | -         | 85            | 11.6 |
| 72                | Severe 78  | F   | CS       | -      | -       |           | TBI    | WAIS-R         | Admission   | -                      | Retro       | 3         | 8.9        | -       | 23.7    | 2.9          | 29.8           | 29.7      | 81            | -    |
| 73                | Severe 71  | PV  | CS       | -      | -       |           | CHI    | WAIS-R         | -           | -                      | -           | 4         | -          | -       | 36.1    | 1.4          | 31.5           | 30.1      | 79            | 11.4 |
| 74                | Severe 30  | FPV | CS       | -      | -       |           | TBI    | WAIS-R         | -           | -                      | -           | 3         | -          | 26.2    | -       | 3.9          | 31.6           | 27.7      | 77            | -    |
| 67                | Severe 28  | F   | CS       | -      | -       |           | TBI    | WAIS-R         | -           | Retro (records)        | -           | 4         | -          | 23.6    | -       | 5.3          | 32.4           | 27.1      | -             | 12.7 |
| 75                | Severe 11  | FPV | CS       | -      | -       |           | HI     | WAIS           | -           | -                      | -           | 2         | -          | 79.8    | -       | 5.4          | 34.7           | 29.3      | 82            | -    |
| 76                | Severe 118 | FPV | CS       | -      | -       |           | CHI    | WAIS-III       | -           | -                      | -           | 4         | 8.4        | -       | 17.3    | 2.9          | 35.5           | 32.6      | 71            | -    |
| 77*               | Severe 24  | F   | CS       | C      | -       |           | CHI    | WAIS-R         | Interview   | -                      | Retro       | 5         | -          | 34.4    | 74.8    | 10.6         | 35.6           | 25.3      | 83            | 14.0 |
| 78                | Severe 35  | F   | CS       | -      | -       |           | TBI    | WAIS-R         | -           | -                      | -           | 2         | -          | 8.9     | 74.5    | 0.7          | 36.1           | 35.4      | 80            | -    |
| 79                | Severe 25  | FPV | CS       | -      | No RTS  |           | TBI    | WAIS-R         | -           | -                      | -           | 3         | -          | 9.8     | -       | 3.1          | 37.0           | 33.9      | 85            | -    |
| 12                | Severe 29  | F   | CS       | -      | -       |           | TBI    | WAIS-III       | Admission   | -                      | -           | 4         | 8.4        | -       | -       | -            | 38.2           | -         | 72            | 12.8 |
| 80                | Severe 76  | FPV | CS       | -      | -       |           | TBI    | WAIS-R         | -           | -                      | -           | 3         | -          | 14      | -       | 14.1         | 38.6           | 24.9      | 83            | 11.9 |
| 81                | Severe 60  | F   | CS       | C      | -       |           | TBI    | WAIS-III       | Lowest 24h  | -                      | Westmead    | 7         | 7.4        | -       | 26.3    | 10.6         | 42.0           | 31.4      | 55            | 12.1 |
| 63                | Severe 16  | F   | CS       | C      | Old     |           | TBI    | WAIS-III       | Admission   | -                      | -           | 4         | 4.7        | -       | 62.5    | 3.8          | 44.4           | 40.6      | 88            | -    |

| Study Methodology         |          |      |          |        |         |           |        |                |             | Injury Characteristics |             |           |                |         | Demographics |         |         |                |           |               |
|---------------------------|----------|------|----------|--------|---------|-----------|--------|----------------|-------------|------------------------|-------------|-----------|----------------|---------|--------------|---------|---------|----------------|-----------|---------------|
| Ref                       | Severity | n    | IQ Index | Design | Control | Subsample | Injury | Wechsler Scale | GCS measure | LOC measure            | PTA measure | NOS (1-9) | GCS (15-3) (d) | LOC (d) | PTA (d)      | TPI (y) | Age (y) | Age Injury (y) | Males (%) | Education (y) |
| Severe                    |          |      |          |        |         |           |        |                |             | Adult Sample           | Mean        | 3.7       | 7.1            | 24.9    | 55.1         | 4.4     | 31.7    | 27.3           | 76.9      | 12.6          |
|                           |          |      |          |        |         |           |        |                |             |                        | SD          | 1.0       | 1.6            | 20.3    | 34.7         | 3.4     | 5.3     | 4.9            | 9.7       | 0.8           |
| Total                     |          | 1766 |          |        |         |           |        |                |             |                        | Mean        | 4.0       | 6.5            | 20.6    | 52.8         | 3.6     | 22.9    | 18.9           | 70.8      | 11.4          |
|                           |          |      |          |        |         |           |        |                |             |                        | SD          | 1.2       | 1.4            | 15.1    | 33.4         | 3.0     | 10.6    | 9.8            | 12.0      | 2.8           |
|                           |          |      |          |        |         |           |        |                |             |                        |             |           |                |         |              |         |         |                |           |               |
| Total TBI severity groups |          | 3116 |          |        |         |           |        |                |             |                        | Mean        | 4.1       | 9.8            | 11.9    | 31.3         | 2.9     | 22.9    | 19.2           | 66.6      | 11.5          |
|                           |          |      |          |        |         |           |        |                |             |                        | SD          | 1.4       | 3.6            | 15.3    | 36.3         | 2.6     | 11.6    | 2.6            | 17.7      | 2.4           |

Note: C = healthy controls; CHI = closed head injury; COAT = Children's Orientation and Amnesia Test; CS = cross-sectional; EI = excellent integration; F = full-scale intelligence quotient; GCS = Glasgow Coma Scale; GOAT = Galveston Orientation and Amnesia Test; HI = head injury; L = longitudinal; LOC = loss of consciousness; MMI = mild memory impairment; Mod = moderate; n = sample size; NOS = Newcastle-Ottawa Scale quality score; OC = orthopedic controls; OGMS = orientation group monitoring system; P= performance intelligence quotient; PI = poor integration; pro = prospective; PTA = post-traumatic amnesia; RE = re-employed; Ref = reference; retro = retrospective; S = severe memory impairment; TOPS = 'TOPS' subgroup; SC = sibling controls; TBI = traumatic brain injury; TPI = time post-injury; TPP= time, place and person; V = verbal intelligence quotient; WAIS = Wechsler Adult Intelligence Scale; WASI = Wechsler Abbreviated Scale for Intelligence; Westmead = Westmead PTA Scale; WISC = Wechsler Intelligence Scale for Children; WPPSI = Wechsler Preschool and Primary Scale of Intelligence.

\*Studies excluded for comparisons between the subacute and chronic phase groups.

## INJURY SEVERITY

### GCS SCORES

The forty-seven studies that reported GCS scores differed in the time of administration of the GCS (also see Table 1). Twenty-five studies measured GCS scores at admission to the hospital. Furthermore, six studies reported the post-resuscitation GCS score, three studies reported the lowest score measured during hospitalization, three studies reported the lowest GCS score measured in the first 24 hours of admission, two studies reported the GCS score at 6 hours and 24 hours post-injury, respectively and one study reported the earliest stable GCS score. Seven studies did not report the time of administration.

### LOC DURATION

Twenty-seven studies reported LOC duration. Of these, eleven studies defined LOC duration as the time between injury and resolution of coma (not further specified). Furthermore, five studies defined LOC duration as the time between injury and the ability to obey simple verbal commands, three studies defined LOC duration as the time between injury and attainment of a GCS score of 15, 9 or 8. Eight studies did not provide a LOC duration definition.

Six studies recorded LOC duration prospectively, of which five studies applied the GCS and one study used clinical observations. Another five studies recorded LOC duration retrospectively, of which four studies used medical records and one study used self-reports to estimate LOC duration. Sixteen studies did not report the methodology of LOC duration measurement.

### PTA DURATION

Twenty-nine studies reported PTA duration. Of these, four studies defined post-traumatic amnesia as the period between regaining consciousness and the recovery of orientation in time, place and person and/or awareness of activities of daily life. In line with the original definitions by Russel & Smith (1961), six studies defined PTA duration as the interval between injury and the recovery of orientation in time, place and person and/or awareness of daily activities. Nineteen of the studies that reported PTA duration did not provide any definition of PTA duration.

Twelve studies reported prospective recordings of PTA duration. Four of these studies used unstandardized clinical judgments of orientation in time, place and person, whereas eight studies applied the standardized Westmead PTA scale, Galveston Orientation and Amnesia Test, Child Orientation and Amnesia Test or Orientation Group Monitoring System. Six studies reported retrospective recordings of PTA duration, of which one estimated PTA



duration using McMillan's structured interview and five studies used hospital notes and self-reports on post-injury orientation and/or amnesia. Eleven studies did not provide any information on the methodology of PTA measurement.

### GROUP COMPARABILITY

To investigate comparability of the TBI severity groups, the mild, moderate and severe TBI groups were compared on age, age at injury, gender ratio, years of education, TPI, GCS score and NOS study quality score (see Table 1). In the subacute phase of recovery, there were no significant differences between the TBI severity groups ( $ts \leq 1.9$ ,  $ps \geq .10$ ), except for longer TPI in the severe group than the mild TBI group ( $t(38) = -2.1$ ,  $p = .048$ ). There were also no significant differences between the TBI severity groups in the chronic phase ( $ts \leq 1.5$ ,  $ps \geq .13$ ), except for longer TPI in the severe TBI group as compared to the mild as well as moderate TBI groups ( $ts \geq 3.5$ ,  $ps \leq .001$ ). In addition, the moderate TBI group had a significant younger age at testing compared to the mild TBI group ( $t(34.0) = 2.1$ ,  $p = .046$ ). A large but non-significant difference in PTA duration was identified between the severe TBI groups in the subacute phase ( $M = 34.7$ ,  $SD = 28.9$ ) and chronic phase ( $M = 52.8$ ,  $SD = 33.4$ ). We matched these groups on injury severity during analyses comparing TPI groups, by excluding four studies from the chronic phase group (references 24, 64, 72-73 in Supporting Information), in order to prevent effects of injury severity to confound the analysis of recovery. As a result of the TBI severity classification procedure, the mild, moderate and severe TBI groups showed progressively lower GCS scores in the in the subacute phase group as well as the chronic phase group ( $ts \geq 5.9$ ,  $ps \leq .001$ ).

As differences in demographic variables between the TPI groups could confound interpretation of recovery effects, we also assessed TPI group comparability. The subacute phase and the chronic phase groups were compared on age, age at injury, gender, years of education, GCS score and NOS study quality score, separately for the mild, moderate and severe TBI groups. For none of the variables of interest a significant difference was obtained between the subacute phase and the chronic phase in the mild TBI group ( $ts \leq 1.6$ ,  $ps \geq .13$ ), the moderate TBI group ( $ts \leq 0.9$ ,  $ps \geq .41$ ) and the severe TBI group ( $ts \leq 1.3$ ,  $ps \geq .20$ ). As a result of the classification procedure for TPI, there was a significant higher TPI in the chronic phase as compared to the subacute phase for all TBI severity groups ( $ts \geq 4.1$ ,  $ps \leq .001$ ).

## PUBLICATION BIAS

Fail-safe *n* values reflecting robustness of effect sizes for the influence of publication bias, indicated that the effect of *mild TBI* on FSIQ in the chronic phase was robust, whereas the effect on VIQ in the chronic phase was not robust (Table 2). Effect sizes in the *moderate TBI* group were not robust to the influence of publication bias according to the fail-safe *n* values. Fail-safe *n* values in the *severe TBI* group indicated robust effect sizes. No funnel plot asymmetry was found in the effect sizes in the mild, moderate or severe TBI groups, except for VIQ of the moderate TBI group in the chronic phase of recovery. There were no associations between the magnitude of effect sizes and sample size or study quality. These results indicate that it is unlikely that publication bias meaningfully influenced the reported effect sizes for the mild, moderate and severe TBI groups, with the single exception of VIQ impairments in the chronic phase after moderate TBI.

## LIST OF ARTICLES INCLUDED FOR META-ANALYSIS

1. Anderson VA, Catroppa C, Rosenfeld JV, Haritou F, Morse SA. Recovery of memory function following traumatic brain injury in pre-school children. *Brain Inj.* 2000;14(8):679–692.
2. Chadwick O, Rutter M, Brown G, Shaffer D, Traub MU. A prospective study of children with head injuries: II. Cognitive sequelae. *Psychol Med.* 1981;11(1):49–61.
3. Levin HS, Song J, Ewing-Cobbs L, Chapman SB, D. M. Word fluency in relation to severity of closed head injury, associated frontal brain lesions, and age at injury in children.pdf. *Neuropsychologia.* 2001;39:122–131.
4. Knights RM, Ivan LP, Ventureya ECG, et al. The effects of head injury in children on neuropsychological and behavioural functioning. *Brain Inj.* 1991;5(4):339–351.
5. Kinsella G, Prior M, Sawyer M, et al. Neuropsychological deficit and academic performance in children and adolescents following traumatic brain injury. *J Pediatr Psychol.* 1995;20(6):753–767.
6. Krivitzky LS, Roebuck-Spencer TM, Roth RM, Blackstone K, Johnson CP, Gioia G. Functional magnetic resonance imaging of working memory and response inhibition in children with mild traumatic brain injury. *J Int Neuropsychol Soc.* 2011;17(6):1143–52.
7. Bassett SS, Slater EJ. Neuropsychological function in adolescents sustaining mild closed head injury. *J Pediatr Psychol.* 1990;15(2):225–236.
8. Shorooei MH, Sharif-Alhoseini M, Saadat S, Sheikh-Mozaffari A, Rahimi-Movaghar V. Effect of mild head injury on intelligence in Zahedan, Iran. *Chinese J Traumatol.* 2010;13(6):345–348.
9. Barth JT, Macciocchi SN, Giordani B, Rimel R, Jane JA, Boll TJ. Neuropsychological sequelae of minor head injury. *Neurosurgery.* 1983;13(5):529–533.
10. Dikmen S, Machamer J, Temkin N. Mild head injury: Facts and artifacts. *J Clin Exp Neuropsychol.* 2001;23(6):729–738.
11. McMillan TM, Glucksman EE. The neuropsychology of moderate head injury. *J Neurol Neurosurg Psychiatry.* 1987;50:393–397.
12. Axelrod BN, Fichtenberg NL, Millis SR, Wertheimer JC. Detecting incomplete effort with Digit Span from the Wechsler Adult Intelligence Scale-third edition. *Clin Neuropsychol.* 2006;20(3):513–523.
13. Kunishio K, Matsumoto Y, Kawada S, et al. Neuropsychological outcome and social recovery of head-injured patients. *Neurol Med Chir.* 1993;33(12):824–829.
14. Tremont G, Mittenberg W, Miller LJ. Acute intellectual effects of pediatric head trauma. *Child Neuropsychol.* 1999;5(2):104–114.
15. Ropacki MT, Elias JW. Preliminary examination of cognitive reserve theory in closed head injury. *Arch Clin Neuropsychol.* 2003;18(6):643–654.
16. Baird A, Papadopoulou K, Greenwood R, Cipolotti L. Memory function after resolution of post-traumatic amnesia. *Brain Inj.* 2005;19(10):811–817.
17. Anderson V A, Moore C. Age at injury as a predictor of outcome following pediatric head injury: a longitudinal perspective. *Child Neuropsychol.* 1995;1(3):187–202.
18. Donders J. Validity of the Kaufman Assessment Battery for Children when employed with children with traumatic brain injury. *J Clin Psychol.* 1992;48(2):225–230.
19. Raghubar KP, Barnes M a, Prasad M, Johnson CP, Ewing-Cobbs L. Mathematical outcomes and working memory in children with TBI and orthopedic injury. *J Int Neuropsychol Soc.* 2013;19(3):1–10. doi:10.1017/S1355617712001312.
20. van Heugten CM, Hendriksen J, Rasquin S, Dijcks B, Jaeken D, Vles JHS. Long-term neuropsychological performance in a cohort of children and adolescents after severe paediatric traumatic brain injury. *Brain Inj.* 2006;20(9):895–903.
21. Hooper SR, Roof KD. Utility of the Hobby WISC-R Split-half Short Form for children and adolescents with severe head injury. *Psychol Rep.* 1993;72(2):371–376.

22. Recla M, Bardoni A, Galbiati S, et al. Cognitive and adaptive functioning after severe TBI in school-aged children. *Brain Inj.* 2013;27(7-8):862–71. doi:10.3109/02699052.2013.775499.
23. Cattelani R, Lombardi F, Brianti R, Mazzucchi A. Traumatic brain injury in childhood: intellectual, behavioural and social outcome into adulthood. *Brain Inj.* 1998;12(4):283–296.
24. Cattelani R, Tanzi F, Lombardi F, Mazzucchi A. Competitive re-employment after severe traumatic brain injury: Clinical, cognitive and behavioural predictive variables. *Brain Inj.* 2002;16(1):51–64.
25. Axelrod BN, Vanderploeg RD, Rawlings DB. WAIS-R prediction equations in patients with traumatic brain injury. *J Clin Exp Neuropsychol.* 1999;21(3):368–374.
26. Oestreicher JM, O'Donnell JP. Validation of the General Neuropsychological Deficit Scale with nondisabled, learning-disabled, and head-injured young adults. *Arch Clin Neuropsychol.* 1995;10(3):185–191.
27. Urakami Y. Relationship between sleep spindles and clinical recovery in patients with traumatic brain injury: A simultaneous EEG and MEG study. *Clin EEG Neurosci.* 2012;43(1):39–47.
28. Watt SE, Shores EA, Baguley IJ, Dorsch N, Fearnside MR. Protein S-100 and neuropsychological functioning following severe traumatic brain injury. *Brain Inj.* 2006;20(10):1007–1017.
29. Kaplan CP, Corrigan JD. Effect of blood alcohol level on recovery from severe closed head injury. *Brain Inj.* 1992;6(4):337–349.
30. Mandleberg IA. Cognitive recovery after severe head injury. 3. WAIS verbal and performance IQs as a function of post-traumatic amnesia duration and time from injury. *J Neurol Neurosurg Psychiatry.* 1976;39(10):1001–1007.
31. Donders J. Psychometric intelligence in patients with traumatic brain injury: utility of a new screening measure. *Arch Phys Med Rehabil.* 1999;80(3):346–347.
32. Kelly MP, Johnson CT, Knoller N, Drubach DA, Winslow MM. Substance abuse, traumatic brain injury and neuropsychological outcome. *Brain Inj.* 1997;11(6):391–402.
33. Catale C, Marique P, Closset A, Meulemans T. Attentional and executive functioning following mild traumatic brain injury in children using the Test for Attentional Performance (TAP) battery. *J Clin Exp Neuropsychol.* 2009;31(3):331–338.
34. Papero PH, Snyder HM, Gotschall CS, Johnson DL, Eichelberger MR. Relationship of two measures of injury severity to pediatric psychological outcome 1-3 years after acute head injury. *J Head Trauma Rehabil.* 1997;12(3):51–67.
35. Max JE, Roberts M a, Koele SL, et al. Cognitive outcome in children and adolescents following severe traumatic brain injury: Influence of psychosocial, psychiatric, and injury-related variables. *J Int Neuropsychol Soc.* 1999;5(1):58–68.
36. Hawley CA, Ward AB, Magnay AR, Mychalkiw W. Return to school after brain injury. *Arch Dis Child.* 2004;89(2):136–142.
37. Levin HS, Eisenberg HM, Wigg NR, Kobayashi K. Memory and intellectual ability after head injury in children and adolescents. *Neurosurgery.* 1982;11(5):668–673.
38. Hall S, Bornstein RA. The relationship between intelligence and memory following minor or mild closed head injury: Greater impairment in memory than intelligence. *J Neurosurg.* 1991;75(3):378–381.
39. Tremont G, Hoffman RG, Scott JG, Adams RL, Nadolne MJ. Clinical utility of Wechsler Memory Scale-Revised and predicted IQ discrepancies in closed head injury. *Arch Clin Neuropsychol.* 1997;12(8):757–762.
40. Skeel RL, Sitzler D, Fogal T, Wells JT, Johnstone B. Comparison of Predicted-difference, Simple-difference, and Premorbid-estimation methodologies for evaluating IQ and memory score discrepancies. *Arch Clin Neuropsychol.* 2004;19(3):363–374.
41. Fisher DC, Ledbetter MF, Cohen NJ, Marmor D, Tulskey DS. WAIS-III and WMS-III profiles of mildly to severely brain-injured patients. *Appl Neuropsychol.* 2000;7(3):126–132.
42. Curtis KL, Greve KW, Bianchini KJ. The Wechsler Adult Intelligence Scale-III and malingering in traumatic brain injury: Classification accuracy in known groups. *Assessment.* 2009;16(4):401–414.
43. Golden Z, Golden CJ. Impact of brain injury severity on personality dysfunction. *Int J Neurosci.* 2003;113(5):733–745.

44. Jeon IC, Kim OL, Kim MS, Kim SH, Chang CH, Bai DS. The effect of premorbid demographic factors on the recovery of neurocognitive function in traumatic brain injury patients. *J Korean Neurosurg Soc.* 2008;44(5):295–302.
45. Dennis M, Agostino A, Roncadin C, Levin H. Theory of mind depends on domain-general executive functions of working memory and cognitive inhibition in children with traumatic brain injury. *J Clin Exp Neuropsychol.* 2009;31(7):835–847.
46. Gale SD, Prigatano GP. Deep white matter volume loss and social reintegration after traumatic brain injury in children. *J Head Trauma Rehabil.* 2010;25(1):15–22.
47. Pentland L, Todd JA, Anderson V. The impact of head injury severity on planning ability in adolescence: A functional analysis. *Neuropsychological.* 1998;8(3):301–317.
48. Wade SL, Walz NC, Carey J, et al. A randomized trial of reen online problem solving for improving executive function deficits following pediatric traumatic brain injury. *J Head Trauma Rehabil.* 2010;25(6):409–415.
49. Wozniak JR, Krach L, Ward E, et al. Neurocognitive and neuroimaging correlates of pediatric traumatic brain injury: a diffusion tensor imaging (DTI) study. *Arch Clin Neuropsychol.* 2007;22(5):555–568.
50. Gensemer IB, Walker JC, McMurry FG, Brotman SJ. IQ levels following trauma. *J Trauma.* 1989;29(12):1616–1619.
51. Kodama T, Morita K, Doi R, Shoji Y, Shigemori M. Neurophysiological analyses in different color environments of cognitive function in patients with traumatic brain injury. *J Neurotrauma.* 2010;27(9):1577–1584.
52. Mathias CW, Greve KW, Bianchini KJ, Houston RJ, Crouch JA. Detecting malingered neurocognitive dysfunction using the reliable digit span in traumatic brain injury. *Assessment.* 2002;9(3):301–308.
53. Roth DL, Conboy TJ, Reeder KP, Boll TJ. Confirmatory factor analysis of the Wechsler Memory Scale-Revised in a sample of head-injured patients. *J Clin Exp Neuropsychol.* 1990;12(6):834–842.
54. Ward H, Shum D, McKinlay L, Baker S, Wallace G. Prospective memory and pediatric traumatic brain injury: Effects of cognitive demand. *Child Neuropsychol.* 2007;13(3):219–39.
55. Souza LMDN, Braga LW, Filho GN, Dellatolas G. Quality-of-life: Child and parent perspectives following severe traumatic brain injury. *Dev Neurorehabil.* 2007;10(1):35–47.
56. Salorio CF, Slomine BS, Grados M, Vasa R, Christensen JR, Gerring JP. Neuroanatomic correlates of CVLT-C performance following pediatric traumatic brain injury. *J Int Neuropsychol Soc.* 2005;11:686–96.
57. Warschausky S, Kewman DG, Selim A. Attentional performance of children with traumatic brain injury: A quantitative and qualitative analysis of digit span. *Arch Clin Neuropsychol.* 1996;11(2):147–153.
58. Mandalis A, Kinsella G, Ong B, Anderson V. Working memory and new learning following pediatric traumatic brain injury. *Dev Neuropsychol.* 2007;32(2):683–701.
59. Perrott SB, Taylor HG, Montes L. Neuropsychological sequelae, familial stress, and environmental adaptation following pediatric head injury. *Dev Neuropsychol.* 1991;7(1):69–86.
60. Park BS, Allen DN, Barney SJ, Ringdahl EN, Mayfield J. Structure of attention in children with traumatic brain injury. *Appl Neuropsychol Adult.* 2009;16(1):1–10. doi:10.1080/09084280802636371.
61. Galbiati S, Recla M, Pastore V, et al. Attention Remediation Following Traumatic Brain Injury in Childhood and Adolescence. *Neuropsychology.* 2009;23(1):40–49.
62. Verger K, Junqué C, Levin HS, et al. Correlation of atrophy measures on MRI with neuropsychological sequelae in children and adolescents with traumatic brain injury. *Brain Inj.* 2001;15(3):211–21. doi:10.1080/02699050010004059.
63. Barker LA, Andrade J, Morton N, Romanowski CAJ, Bowles DP. Investigating the “latent” deficit hypothesis: age at time of head injury, implicit and executive functions and behavioral insight. *Neuropsychologia.* 2010;48:2550–2563.
64. Ferri J, Chirivella J, Caballero MC, Simo M, Ramirez RM, Noe E. Intelligence and prognosis in severe traumatic brain injury: A neuropsychological study with the Wechsler adult intelligence scale (WAIS-III). *Neurologia.* 2004;19(6):285–291.

65. Goldstein FC, Levin HS, Boake C. Conceptual encoding following severe closed head injury. *Cortex*. 1989;25(4):541–554.
66. Burke JM, Imhoff CL, Kerrigan JM. MMPI correlates among post-acute TBI patients. *Brain Inj*. 1990;4(3):223–231.
67. Goran DA, Fabiano RJ, Crewe N. Employment following severe traumatic brain injury: The utility of the Individual Ability Profile system (IAP). *Arch Clin Neuropsychol*. 1997;12(7):691–698.
68. Johnson SC, Bigler ED, Burr RB, Blatter DD. White matter atrophy, ventricular dilation, and intellectual functioning following traumatic brain injury. *Neuropsychology*. 1994;8(3):307–315.
69. Schmitter-Edgecombe ME, Marks W, Fahy JF, Long CJ. Effects of severe closed-head injury on three stages of information processing. *J Clin Exp Neuropsychol*. 1992;14(5):717–737.
70. Trudel TM, Tryon WW, Purdum CM. Awareness of disability and long-term outcome after traumatic brain injury. *Rehabil Psychol*. 1998;43(4):267–281.
71. Fleming J, Riley L, Gill H, Gullo MJ, Strong J, Shum D. Predictors of prospective memory in adults with traumatic brain injury. *J Int Neuropsychol Soc*. 2008;14(5):823–31.
72. Weddell RA. Relatives' criticism influences adjustment and outcome after traumatic brain injury. *Arch Phys Med Rehabil*. 2010;91(6):897–904.
73. Paniak CE, Silver K, Finlayson MAJ, Tuff LP. How useful is the WAIS-R in closed head injury assessment? *J Clin Psychol*. 1992;48(2):219–225.
74. Reza F, Ikoma K, Chuma T, Mano Y. Correlations between neuropsychological test results and P300 latency during silent-count and button-press tasks in post-traumatic brain injury patients. *J Clin Neurosci*. 2006;13(9):917–922.
75. Newton A, Johnson DA. Social adjustment and interaction after severe head injury. *Br J Clin Psychol*. 1985;24(4):225–234.
76. Wood RLL, Liossi C. The relationship between general intellectual ability and performance on ecologically valid executive tests in a severe brain injury sample. *J Int Neuropsychol Soc*. 2007;13(1):90–98.
77. Schmitter-Edgecombe M, Marks W, Wright MJ, Ventura M. Retrieval inhibition in directed forgetting following severe closed-head injury. *Neuropsychology*. 2004;18(1):104–14.
78. Baek SO, Kim OL, Kim SH, et al. Relation between cingulum injury and cognition in chronic patients with traumatic brain injury; diffusion tensor tractography study. *NeuroRehabilitation*. 2013;33(3):465–71. doi:10.3233/NRE-130979.
79. Dornan J, Schentag C. Traumatic brain injury: factors predicting return to work or school. *Brain Inj*. 1995;9(5):517–532.
80. Hoofien D, Gilboa A, Vakil E, Donovick PJ. Traumatic brain injury (TBI) 10-20 years later: a comprehensive outcome study of psychiatric symptomatology, cognitive abilities and psychosocial functioning. *Brain Inj*. 2001;15(3):189–209.
81. Draper K, Ponsford J. Cognitive functioning ten years following traumatic brain injury and rehabilitation. *Neuropsychology*. 2008;22(5):618–625.